



LIQUID FERTILIZER

MOLY 6

TO BE USED WHERE A MOLYBDENUM DEFICIENCY EXISTS.

GUARANTEED MINIMUM ANALYSIS
SODIUM MOLYBDATE AS NATURAL PHENOLIC COMPLEX 6% W/W
BORON 0.4% W/W

5 Litre NET **20 Litre NET** **200 Litre NET**

25 BS-2

Moly 6 application recommendations: Apply to the following crops where a deficiency exists.

Minimum water rate is 1:50 for foliar application and 1:100 for fertigation.

CROP	FOLIAR RATES L/ha	FERTIGATION RATES L/ha	COMMENTS
ALMONDS	1-2L	3-6L	For best results apply onto new foliage when symptoms are exposed.
APPLES	1-1.5L	3-6L	For best results apply onto new foliage when symptoms are exposed.
ASPARAGUS	0.5-1L	2-4L	Apply as symptoms are exposed.
AVOCADOS	1-3L	4-8L	Apply as symptoms are exposed or apply at 4-5 leaf stage.
BANANAS	0.5-1.5L	2-4L	Apply as symptoms are exposed.
BROCCOLI	0.5-1L	2-4L	Apply as symptoms are exposed or apply at 4-5 leaf stage.
CARPSICUM	0.5-1L	2-4L	Apply as symptoms are exposed or apply at 4-5 leaf stage.
CELERY	0.2-1.2L	2-4L	Apply as symptoms are exposed or apply at 4-5 leaf stage.
CITRUS	1-1.5L	4-7L	For best results apply onto new foliage when symptoms are exposed.
COTTON	0.6-1.5L	4-8L	For best results apply onto new foliage when symptoms are exposed.
CUCUMBERS	1-2L	3-6L	Apply as symptoms are exposed or apply at 4-5 leaf stage.
GRAPE VINES	1-2L	3-6L	For best results apply onto new foliage when symptoms are exposed.
LEGUMES	1-2L	3-7L	Apply when early growth is approx. 150mm tall to ensure effective movement into nitrogen fixing nodules.
MELONS	0.5-1L	2-4L	Apply as symptoms are exposed or apply at 4-5 leaf stage.
NUT CROPS	0.3-1L	2-3L	For best results apply onto new foliage when symptoms are exposed.

CROP	FOLIAR RATES L/ha	FERTIGATION RATES L/ha	COMMENTS
OLIVES	0.3-1L	2-3L	For best results apply onto new foliage when symptoms are exposed.
PASTURE/LUCERNE	0.5-1L	1-2L	Apply when early growth is approx. 150mm tall to ensure effective movement into nitrogen fixing nodules. Apply at least 2 weeks before cutting.
POTATOES	0.4-1L	1-2L	Apply as symptoms are exposed or apply at 4-5 leaf stage.
STONE FRUIT	0.3-1L	1-2L	For best results apply onto new foliage when symptoms are exposed.
STRAWBERRIES	0.5-1L	1.5-2.5L	Apply as symptoms are exposed or apply at 4-5 leaf stage.
TOMATOES	0.4-1L	1-3L	Apply as symptoms are exposed or apply at 4-5 leaf stage.
TREE CROPS	1-2L	3-6L	For best results apply onto new foliage when symptoms are exposed.
VEGETABLES	0.5-1L	1.5-3L	Apply as symptoms are exposed or apply at 4-5 leaf stage.
ZUCCHINI	0.4-1L	1-2L	Apply as symptoms are exposed or apply at 4-5 leaf stage.

ANALYSIS	AUSTRALIA W/V %	INTERNATIONAL W/V %
Molybdenum	6.0	5.3
Boron	0.4	0.35

Specific gravity 1.10 pH 7-9



CAUTION: This product is harmful if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Avoid breathing spray mist.

*Warning: Excessive use of molybdenum can be harmful to stock. Do not apply fertilizer containing molybdenum on more than 25% of farm annually. Plant levels of molybdenum can be high for 4 weeks after application. It is advisable to keep stock off treated areas for this period. Check rate and frequency of molybdenum use with appropriate authorities. The rates must be adjusted according to the tissue and/or soil analysis results. **STORAGE & DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinse into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

NOT APPLICABLE TO 1000 LITRE CONTAINERS

SPRAYGRO LIQUID FERTILIZERS
 76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE PORT ADELAIDE, SOUTH AUSTRALIA 5015
 PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@spraygro.com.au • www.spraygro.com.au • A.B.N. 47 007 974 498

PRODUCED AND MANUFACTURED IN AUSTRALIA

CAUTION

The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture.
- Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day.
- **Physically compatible** with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.
- Use the correct label spray rates of any pesticide used with SprayGro product.
- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day.

NOTE: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

Safety glasses, gloves and protective clothing should be worn when using this product.

DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizer and its servants and agents ("Spraygro") shall be under no liability to the purchaser and/or user of Spraygro products ("customer") in respect of any loss or damage (including consequential loss or damage which may be suffered, incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon and descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer must independently verify the accuracy thereof. 4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability reasonably incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our Distributor.



BATCH No.