



Suggested Nutrient Applications for:



Apples & Pears

Latest Update: 18/11/10

Growth Stage	Silver Tip	Green Tip	Pink Bud	Start of Flowering	Petal Fall	End of Flowering	Fruit Set	Fruit Development	2 Week Harvest	Before Leaf Senescence
Product % Nutrient (N-P-K) w/v										
Humic Acid 12 12 HA	15									
Base™ 15-18-20 + 0.1 Fe, 0.08 Mn, 0.1 Zn, 0.04Cu, 0.03B, 0.022Mg, 0.002Mo	30								10	
Nitro 35 35 - 0 - 0		15				10		10		
Smartrace® Zinc 10 Zn, 4.9 S		2								5
Firmrite® CB 2 B, 16 Ca			2	2	10		10			
Smartrace® Spraytrace 8 Fe, Zn, Mg, Mn, S, B, N, Cu, Mo				5					6	
Complete K 5 - 10 - 27, 0.02 Mg, Zn, Mn, Cu, Fe, B, 0.002 Mo							5		5	
Smartrace® Calcium 3.5 - 0 - 0, 5 Ca							4	4	4	
Firmrite® CMB 15-0-0 + 22 Ca, 3 Mg, 0.1 B					5		5			

**Rate
Litres/Hectare**

= PRODUCT TO PRODUCT COMPATIBILITY*

* = Product to Product Compatibility when diluted with water at correct label rates

Minimum Water Rate: Foliar Application 1:50 Fertigation 1:100

= Foliar (Lt/Ha)

= Fertigation (Lt/Ha)

Important Notes:

- (1) The product label should be consulted before use of any of the products referred to on this sheet, always refer to the label when applying SprayGro Liquid Fertilizer products.
- (2) To ensure correct nutrients are delivered to the plant, tissue testing should be carried out prior applying any nutrients to determine your crop situation. Crop damage is possible if trace elements are applied in excess more so than from deficiency.
- (3) This information sheet is a guide only as SprayGro Liquid Fertilizers neither makes, nor authorizes any agent or representative to make any other warranties, express or implied, other than those stated on the product label.
- (4) SprayGro Liquid Fertilizers shall not be liable for any results, loss, or damage whatsoever, whether consequential or otherwise through the use or application of products and/or materials referred to herein.
- (5) We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray. Combination sprays can induce crop phytotoxicity.
- (6) Apply at times of low light intensity, early morning or preferably during the evening. Avoid applications to wet leaves, stressed crops and on exceptionally hot, dry or sunny days. Overcast, calm and cool conditions are best.
- (7) E.&O.E.