

**DISCLAIMER/CONDITIONS OF SALE:** 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizer and its servants and agents ("Spraygro") shall be under no liability to the purchaser and/or user of Spraygro products ("customer") in respect of any loss or damage including consequential loss or damage which may be suffered, incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon and descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer must independently verify the accuracy thereof. 4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability reasonably incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our Distributor.

BATCH No.



**Safety glasses, gloves and protective clothing should be worn when using this product.**

**NOTE:** We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

- **DO NOT** apply during the heat of the day.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.
- Use the correct label spray rates of any pesticide used with Spraygro product.
- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
- test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- **Physically compatible** with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity (e.g. bucket
- Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day.
- Use Spraygro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture, to the plant.

The following conditions must be observed in order to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage

**CAUTION**



# Gripaper™

edta chelate

## C-TRACE

3.0% Zn, 2% Mn, 1.3% Fe, 0.7% B, 0.4% Mo, 0.3% Cu

20 Litre NETO 200 Litre NETO 1000 Litre NETO

Apply C Trace when crop trace element deficiencies exist.

CROP	RATE/ha	MINIMUM DILUTION	COMMENTS
ALMONDS Foliar Fertigation	2-4L 8-10L	1:100 1:100	Apply as required to maintain crop colour and vigour. Repeat early fruit growth.
CAPSICUM Foliar Fertigation	2-4L 5-10L	1:100 1:100	Apply when sufficient leaf area is present. Repeat at fruit set and development.
CITRUS Foliar Fertigation	4-5L 10-15L	1:100 1:100	Apply as required to maintain trace element levels.
COTTON Aerial	3-7L	Use maximum practical water 1:100	Apply as required to maintain trace element levels.
Ground rig	3-7L		
GRAPES/ VINES Foliar Fertigation	3-5L 10-12L	1:100 1:100	Maximum 3 applications may be required post flowering up to veraison.
OLIVES Foliar Fertigation	3-6L 9-15L	1:100 1:100	Apply early in the season, repeat to maintain trace elements levels. Repeat post-harvest.
PISTACHIOS Foliar Fertigation	2-5L 10-15L	1:100 1:100	Apply as required to maintain crop colour and vigour. Repeat mid-late fruit growth.
POTATOES Foliar Fertigation	3-5L 9-12L	1:100 1:100	Apply to vines when sufficient leaf area. Repeat at tubor development.

CROP	RATE/ha	MINIMUM DILUTION	COMMENTS
STONE FRUIT Fertigation only	10-15L	1:100	Apply pre-flowering or after recently hardened flush. Repeat as necessary.
STRAWBERRIES Foliar Fertigation	2-4L 6-10L	1:100 1:100	Apply 3-4 weeks after transplanting. Repeat at fruit development.
SUGARCANE Foliar Fertigation	4-5L 4-5L	1:100 1:100	Apply during early growth when sufficient leaf area. Apply in conjunction with blight fungicide.
TOMATOES Foliar Fertigation	4-6L 10-15L	1:100 1:100	Apply 3 weeks after transplanting. Repeat at first truss and fruit development.
TURF Foliar Fertigation	3L 6L	1:100 1:100	As required to maintain turf quality and boost trace element levels.

ANALYSIS	AUSTRALIA W/W %	INTERNATIONAL W/W %
Zinc (Zn) as EDTA	3.0	2.5
Manganese (Mn) as EDTA	2.0	1.66
Iron (Fe) as EDTA	1.3	1.08
Copper (Cu) as EDTA	0.3	0.25
Boron (B)	0.7	0.58
Molybdenum (Mo)	0.4	0.33
Sulphur (S)	2.8	2.33

Specific gravity 1.19

**WATER RATIO:** For foliar application use 1:100 minimum water dilution rates. For aerial application use maximum practical water rates. The rates must be adjusted according to the tissue and/or soil analysis results. **NOTE:** Only mix diluted chelate formulations with liquid NPK's and do not store solution.

**STORAGE & DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.



**SPRAYGRO LIQUID FERTILIZERS**

76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE  
PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 996

PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@spraygro.com.au • www.spraygro.com.au

**PRODUCED AND MANUFACTURED IN AUSTRALIA**

NOT APPLICABLE TO 1000 LITRE CONTAINERS