



20 Litre NET 200 Litre NET 1000 Litre NET

Application recommendations: Gripper Copper is suitable for the following crops where Copper deficiency exists. Apply with sufficient water for thorough leaf coverage.

CROP	RATE/ha	MINIMUM DILUTION	COMMENTS
ALMONDS NUT TREES	1-2L 4-6L	1:100 1:100	Apply as required to maintain crop colour and vigour. Repeat early fruit growth.
	0.5-2L	1:40	Apply to actively growing plants when sufficient leaf area exists to spray. Repeat as necessary. Apply between mid-late tillering to flag leaf emergence (fle) stage. After fle apply maximum of 1.0L/ha with minimum of 75L water dilution. Apply at 4-8 leaf stage. Apply prior pod set.
BROADACRE CROPS Cereals Canola Legumes	1-2L 2-4L	1:100 1:100	Apply when sufficient leaf area is present before flowering.
CITRUS	Foliar Fertigation	1:100 1:100	Apply when deficient. Use lower rates on trees up to 6 year stage.
	Aerial	0.89-2L	Use maximum practical water
GRAPES/ VINES	Ground rig Foliar Fertigation	1:50 0.5-1L 2-3L	Apply when deficiency exists.

CROP	RATE/ha	MINIMUM DILUTION	COMMENTS
ONIONS	Foliar Fertigation	1L 1:100	Apply at 2-3 leaf stage Apply at bulb formation.
	Foliar	0.5-2L	Apply in Spring up until bloom. Do not apply during blossom or harvest.
POME FRUIT	Foliar Fertigation	4-6L 1:300	Apply to vines when sufficient leaf area. Repeat at tuber development.
	Foliar	1-2L 1:100	Apply pre-flowering or on recently hardened flush. Repeat as necessary.
STONE FRUIT	Foliar only	3-5L 1:100	Apply 3-4 weeks after transplanting. Repeat at fruit development.
	Foliar	0.5L-1L 1-2L	Apply 3 weeks after transplanting. Repeat at first truss and fruit development.
TOMATOES	Foliar Fertigation	2-3L 4-5L	As required to maintain turf quality and improves colour.
	Foliar	100mls per 100m ²	

ANALYSIS	AUSTRALIA W/V %	INTERNATIONAL W/V %
Copper (Cu) as EDTA	6.0	4.8

Specific Gravity 1.24



WARNING: DO NOT FOLIAR APPLY TO COPPER SENSITIVE TREES SUCH AS APRICOTS, NECTARINES AND PLUMS.
WATER RATIO: For foliar application use 1:100 minimum water dilution rates. For aerial application use maximum practical water rates. The rates must be adjusted according to the tissue and/or soil analysis results. **NOTE: Only mix diluted chelate formulations with liquid NPK's and do not store solution.**

STORAGE & DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

SPRAYGRO LIQUID FERTILIZERS
 76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494, PORT ADELAIDE BUSINESS CENTRE
 PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
 PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@spraygro.com.au • www.spraygro.com.au

PRODUCED AND MANUFACTURED IN AUSTRALIA

NOT APPLICABLE TO 1000 LITRE CONTAINERS

CAUTION


The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture.
- **Early morning or late afternoon** spraying is most effective as evaporation can occur during hotter parts of the day.
- **Physically compatible** with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.
- Use the correct label spray rates of any pesticide used with SprayGro product.
- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day.

NOTE: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

Safety glasses, gloves and protective clothing should be worn when using this product.

DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizer and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect of any loss or damage including consequential loss or damage which may be suffered, incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon and descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer must independently verify the accuracy thereof. 4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability reasonably incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised office of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our Distributor.



BATCH No.