


BATCH No.



DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizer and its servants and agents ("Spraygro") shall be under no liability to the purchaser and/or user of Spraygro products ("customer") in respect of any loss or damage including consequential loss or damage which may be suffered, incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon and descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer must independently verify the accuracy thereof. 4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability reasonably incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our Distributor.

Safety glasses, gloves and protective clothing should be worn when using this product.

and observed prior to the total crop spray.

NOTE: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed

• **DO NOT** apply during the heat of the day.

• **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.

• **DO NOT** apply during the heat of the day.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.

• **DO NOT** apply during the heat of the day.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

Gripper™

edta chelate

ZINC COPPER

8.7 N, 6 Zn, 3 Cu, % w/v

20 Litre NET 200 Litre NET 1000 Litre NET

CAUTION

The following conditions must be observed in order to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.

• **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.



Application recommendations: Gripper Zinc Copper is suitable for the following crops where Zinc Copper deficiency exists. Apply with sufficient water for thorough leaf coverage.

CROP	RATE/ha	MINIMUM DILUTION	COMMENTS
ALMONDS	Foliar 2-3L Fertigation 6-10L	1:100	Apply as required to maintain crop colour and vigour. Repeat early fruit growth.
NUT TREES	Foliar 1-4L	1:40	Apply to actively growing plants when sufficient leaf area exists to spray. Repeat as necessary. Apply between mid-late flowering to flag leaf emergence (file) stage. After file apply maximum of 1.0L/ha with minimum of 75L water dilution. Apply at 4-8 leaf stage. Apply prior pod set.
BROADACRE Cereals Canola Legumes	Foliar 1-3L Foliar 4-8L	1:100	Apply when deficient. Use lower rates on trees up to 6 year stage.
VEGETABLES	Foliar 2-4L Fertigation 6-10L	1:100	Apply when sufficient leaf area is present before flowering.
CITRUS	Foliar 1-3L Fertigation 4-8L	1:100	Apply when deficient. Use lower rates on trees up to 6 year stage.
COTTON	Aerial 1.6-4L	Use maximum practical water	Apply prior squaring.
GRAPES/ VINES	Ground rig 1-2L Foliar 4-8L	1:50 1:100	Apply when deficiency exists.

CROP	RATE/ha	MINIMUM DILUTION	COMMENTS
ONIONS	Foliar 2L Fertigation 4L	1:100	Apply at 2-3 leaf stage Apply at bulb formation.
POME FRUIT	Fertigation 6-10L only	1:300	Fornightly applications during growing season as required, subject to leaf analysis tests/results.
POTATOES	Foliar 3-5L Fertigation 8-10L	1:100	Apply to vines when sufficient leaf area. Repeat at tuber development.
STONE FRUIT	Fertigation 6-10L only	1:100	Apply pre-flowering or on recently hardened flush. Repeat as necessary.
STRAWBERRIES	Foliar 1-2L Fertigation 2-4L	1:100	Apply 3-4 weeks after transplanting. Repeat at fruit development.
TOMATOES	Foliar 2-4L Fertigation 6-10L	1:100	Apply 3 weeks after transplanting. Repeat at first truss and fruit development.
POPPIES	Foliar 3-4 Fertigation 6-10	1:100	Apply from row cover until petal drop as required.

ANALYSIS	AUSTRALIA W/V %	INTERNATIONAL W/W %
Copper (Cu) as EDTA	3.0	2.4
Zinc (Zn) as EDTA	6.0	4.8

Specific Gravity 1.23 ± 0.02

WARNING: DO NOT FOLIAR APPLY TO COPPER SENSITIVE TREES SUCH AS APRICOTS, NECTARINES AND PLUMS.
WATER RATIO: For foliar application use 1:100 minimum water dilution rates. For aerial application use maximum practical water rates. The rates must be adjusted according to the tissue and/or soil analysis results. **NOTE:** Only mix diluted chelate formulations with liquid NPK's and do not store solution.
STORAGE & DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Triple rinse container empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.



NOT APPLICABLE TO 1000 LITRE CONTAINERS

SPRAYGRO LIQUID FERTILIZERS
 76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE, PORT ADELAIDE, SOUTH AUSTRALIA 5015
 A.B.N. 47 007 974 496 • PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@spraygro.com.au • www.spraygro.com.au

PRODUCED AND MANUFACTURED IN AUSTRALIA