


BATCH NO.



DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizer and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect of any loss or damage including consequential loss or damage which may be suffered, incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon and descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer must independently verify the accuracy thereof. 4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability reasonably incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our Distributor.

Safety glasses, gloves and protective clothing should be worn when using this product.

NOTE: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day.

- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.
- **DO NOT** use the correct label spray rates of any pesticide used with SprayGro product.
- **DO NOT** use when precipitation (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.
- Use the correct label spray rates of any pesticide used with SprayGro product.
- **Physically compatible** with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.

- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day.
- **Physically compatible** with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.

The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in damage to the plant.

CAUTION

85 GR-3



Gripper™
edta chelate
7.5% ZINC w/v

20 Litre NET 200 Litre NET 1000 Litre NET



Application recommendations: Spray Gro Gripper Zinc is suitable for the following crops where a zinc deficiency exists. Apply with sufficient water for thorough leaf coverage.

CROP	RATE/ha	MINIMUM DILUTION	COMMENTS
ALMONDS	Foliar Fertigation 1-2L 6-8L	1:100 1:100	Apply as required to maintain crop colour and vigour. Repeat early kernel growth.
BROADACRE CROPS Cereals, Canola Foliar Fertigation Legumes	2-3L 4-6L	1:40 1:100	Apply from early emergence up to tillering. Apply from early emergence where there is high zinc requirement.
CAPSICUM	Foliar Fertigation 2L 4-6L	1:100 1:100	Apply when sufficient leaf area is present. Repeat at fruit set and development.
CITRUS	Foliar Fertigation 2-3L 6-10L	1:100 1:100	Apply as required to maintain trace element levels.
COTTON	Aerial 2L	Use maximum practical water 1:100	Apply as required to maintain trace element levels.
GRAPES/ VINES	Foliar Fertigation 2-3L 4-8L	1:100 1:100	Apply at shoots 10cm, 5% flower and post harvest.
OLIVES	Foliar Fertigation 2-3L 6-10L	1:100 1:100	Apply early in the season, repeat to maintain trace elements levels. Repeat post-harvest.

CROP	RATE/ha	MINIMUM DILUTION	COMMENTS
PISTACHIOS	Foliar Fertigation 1-2L 6-10L	1:100 1:100	Apply as required to maintain crop colour and vigour. Repeat mid-late fruit growth.
POTATOES	Foliar Fertigation 2-3L 6-10L	1:100 1:100	Apply to tops when sufficient leaf area. Repeat at tubor development.
STONE FRUIT	Fertigation 6-10L	1:100	Apply pre-flowering or on recently hardened flush. Repeat as necessary.
STRAWBERRIES	Foliar Fertigation 2L 4-6L	1:50 1:100	Apply 3-4 weeks after transplanting. Repeat as required.
SUGARCANE	Foliar Fertigation 2-3L 4-5L	1:100 1:100	Apply during early growth when sufficient leaf area. Apply in conjunction with billet fungicide.
TOMATOES	Foliar Fertigation 2L 6-10L	1:100 1:100	Apply 3 weeks after transplanting. Repeat at first truss and fruit development.
TURF	Foliar 20L	1:100	As required to maintain turf quality.
VEGETABLES	Foliar Fertigation 2-4L 4-6L	1:100 1:100	Apply at early stage of growth as required.

ANALYSIS	AUSTRALIA W/V %	INTERNATIONAL W/W %
Zinc (Zn) as EDTA	7.5	6.0

Specific gravity 1.24

WATER RATIO: A dilution of 1:100 means 1part product : 100 parts water. In hot weather, use the higher dilution rate when applicable. **NOTE:** Only mix diluted chelate formulation with liquid NPD's and do not store solution. For aerial application use maximum practical water rates. The rates must be adjusted according to the tissue and/or soil analysis results. For optimal results ensure spray tank pH is below 6.

STORAGE & DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

SPRAYGRO LIQUID FERTILIZERS 76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494
PORT ADELAIDE BUSINESS CENTRE, PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@spraygro.com.au • www.spraygro.com.au



NOT APPLICABLE TO 1000 LITRE CONTAINERS

PRODUCED AND MANUFACTURED IN AUSTRALIA