



# Suggested Nutrient Applications for:



Latest Update: 24/11/10

Growth Stage		Autumn	Shoot Growth	Bud Swelling	Flowering	Fruit set	Early Fruiting	Pit Hardening	Fruit Expansion	Post Harvest	Notes
Product	% Nutrient (N-P-K) w/v										
BASE™	15 – 18 – 20 + TE + 0.3 Ha	30		50		50	<b>Rate Litres/Hectare</b>			30	Improves root and shoot growth by providing balanced nutrition; fortified with APP and EDTA trace elements.
FIRMRITE CMB	15 –0 –0 22 Ca, 3 Mg, 0.1 B	5	30		30		30		5	5	Removes calcium related disorders, improves fruit wall and minimize fruit rot.
BORON-15	6 - 0 – 0 + 15 B			1 ←				1 ←	500 ml ↑		Boron improves flowering and fruit set Eliminates soft nose in Kalamata.
SMARTRACE® Zinc	10 Zn, 4.9 S			5 ←		5 ←					Zinc increase IAA production required for shoot growth and to improve fruit size.
PICK® 15-42	0 – 15 – 42	↔ = PRODUCT TO PRODUCT COMPATIBILITY*					5	5	5 ←		Phosphorous and potassium improves oil biosynthesis and fruit size.
SMARTRACE® Spraytrace 8	0.2 – 0 – 0 + 1.3 Zn, 1.3 Fe, 1 Mn, 0.2 Cu, 0.5 B, 0.2 Mo, 2 S		5			5 ←		5 ←		5	Multinutrient lignosulfonate chelated trace elements to remove deficiencies and restore growth.

\* = Product to Product Compatibility when first diluted with water at correct label rates

Minimum Water Rate: Foliar Application 1:50 Fertiligation 1:500



= Foliar (Lt/Ha)



= Fertiligation (Lt/Ha)

### Important Notes:

- (1) The product label should be consulted before use of any of the products referred to on this sheet, always refer to the label when applying SprayGro Liquid Fertilizer products.
- (2) To ensure correct nutrients are delivered to the plant, tissue testing should be carried out prior applying any nutrients to determine your crop situation. Crop damage is possible if trace elements are applied in excess more so than from deficiency.
- (3) This information sheet is a guide only as SprayGro Liquid Fertilizers neither makes, nor authorizes any agent or representative to make any other warranties, express or implied, other than those stated on the product label.
- (4) SprayGro Liquid Fertilizers shall not be liable for any results, loss, or damage whatsoever, whether consequential or otherwise through the use or application of products and/or materials referred to herein.
- (5) We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray. Combination sprays can induce crop phytotoxicity.
- (6) Apply at times of low light intensity, early morning or preferably during the evening. Avoid applications to wet leaves, stressed crops and on exceptionally hot, dry or sunny days. Overcast, calm and cool conditions are best.
- (7) E.&O.E.