

SPRAYGRO®

SMARTTRACE® CHELATED ELEMENT

MANGANESE COPPER

GUARANTEED MINIMUM ANALYSIS

MANGANESE 8% W/V PLUS COPPER 1.5% W/V AS NATURAL PHENOLIC COMPLEX

20 Litre NET • 200 Litre NET • 1000 Litre NET •

APPLICATION RECOMMENDATIONS: SPRAYGRO SMARTTRACE MANGANESE COPPER is suitable for the following crops where a Manganese Copper deficiency exists. Apply with sufficient water for thorough coverage of leaves.

BROADACRE:

Rate of application per min water/hectare: 50 litres

@ 4-6 litres/hectare. Apply when deficiency is known to exist.

@ 4-6 litres/hectare. Apply between full-bud and the onset of flowering.

@ 4-6 litres/hectare. Apply to actively growing plants when sufficient leaf area exists to spray. Repeat as necessary.

Apply mid-late tillering, before early boot stage.

@ 4-6 litres/hectare. Apply 4-6 weeks after planting or when deficiency symptoms appear.

@ 4-6 litres/hectare. Apply at flowering of 1st laterals.

HORTICULTURE:

Rate of application per min water/hectare: 300 litres

CITRUS, POME FRUIT, STONE FRUIT

@ 3-5 litres/hectare. Apply in springtime as a thorough spray, up until bloom and again after bloom if necessary. Repeat applications may be necessary for a severe deficiency. Best absorption occurs 1/2 – 1/3 through a flush of growth. Spray should not be applied during blossoming or harvest.

@ 2-3 litres/hectare. Apply when deficiency symptoms appear.

@ 2-4 litres/hectare. Apply as a thorough cover spray 2-3 weeks before bloom. Do not apply during bloom. Repeat as necessary.

@ 2-4 litres/hectare. Apply a week after transplanting then again prior to bloom if necessary. Young foliage, flowers or fruit are susceptible to trace-element spray damage. Test a few plants or a plot before application.

@ 3-6 litres/hectare. Apply when plants are young and before flowering. Repeat as necessary.

Turf:

Rate of application per min water/hectare: 300 litres

@ 100mls/100²water dilution 1:100.

CAUTION

The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day.
- **Compatible** with a wide range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.
- Use the correct label spray rates of any pesticide used with SprayGro product.
- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day.

NOTE: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

Safety glasses, gloves and protective clothing should be worn when using this product.

CONDITIONS OF SALE: 1) Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than the officer of the Seller, is authorised to make any warranty, guarantee or direction concerning this product. 2) Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

SPRAYGRO LIQUID FERTILIZERS PTY. LTD.

52 BEDFORD STREET, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE
PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@picknowl.com.au

BATCH No.

PRODUCED AND MANUFACTURED IN AUSTRALIA