



# Suggested Nutrient Applications for:

# Strawberries

Latest Update: 24/11/10

Growth Stage		Transplant	7-14 days post plant	14-21 days post plant	Green Bud	White Bud	Flowering	Fruit Set	Fruit Fill	Notes
Product	% Nutrient (N-P-K) W/V									
Humic Acid 12	12 Humic Acid	15			10			10		Improves soil structure, reduces nutrient leaching and improves microbial activity.
Reach	Natural Rooting Hormone & Biostimulant	2		2						Eliminates transplant shock & drives early root growth
Companion®	Concentrated Microbial Inoculant	Dip/Runner 500ml/400 litres water	2			2				Promotes ecological balance thus suppressing pathogen attack.
Base	15 – 18 – 20 + 0.1 S, 0.1 Zn, 0.1 Mn, 0.1 Fe, 0.02 Mg, 0.03 B, 0.04 Cu, 0.002 Mo, 0.3 HA	15		20						Complete starter, ideal for early growth & root establishment. Excellent in ratoon crops
Sprayphos 620	620g/L Phosphorous acid	1.94	1.94	1.94	1.94					Apply for the control of Crown Rot (Phytophthora spp)
Nitro 35	35 – 0 – 0		20		20				3	Long lasting N, reduced leaching and low salt index.
Boron 15	6 – 0 – 0 + 15 B			0.5		1.5		1.5	1.5	Non sodium form of Boron, which drives Calcium uptake, improves flower pollination, set & quality.
Firmrite Spray Cat 7.5	7.5 Ca, 12.5 S			4		4		4	4	Thiosulphate Calcium for better uptake and transport in the plant.
Gripper Magnesium	6.2 – 0 – 0 + 5 Mg			3		3		3	3	EDTA Chelated Magnesium for better uptake and sugar movement
KELPIT	100% Seaweed Sargussam & Durvillea		10		5		5			Seaweed extract supplemented with long lasting Nitrogen and natural Cytokinins
Gripper C Trace	3.4 S, 3 Zn, 2 Mn, 1.3 Fe, 0.7 B, 0.4 Mo, 0.3 Cu				4		4		4	EDTA Chelated multi-trace elements product in an ideal ratio for strawberries.
K Blast 36	0 – 0 – 30 + 24S				6		6		6	High analysis potassium with good uptake

**Litres/Hectare Rate**

**= Product to Product Compatibility**

\* = Product to Product Compatibility when diluted with water at correct rates

Minimum Water Rate: Foliar Application 1:50 Fertigation 1:100  = Foliar (Lt/Ha)  = Fertigation(Lt/Ha)

**Important Notes:**

- (1) The product label should be consulted before use of any of the products referred to on this sheet, always refer to the label when applying SprayGro Liquid Fertilizer products.
- (2) To ensure correct nutrients are delivered to the plant, tissue testing should be carried out prior applying any nutrients to determine your crop situation. Crop damage is possible if trace elements are applied in excess more so than from deficiency.
- (3) This information sheet is a guide only as SprayGro Liquid Fertilizers neither makes, nor authorizes any agent or representative to make any other warranties, express or implied, other than those stated on the product label.
- (4) SprayGro Liquid Fertilizers shall not be liable for any results, loss, or damage whatsoever, whether consequential or otherwise through the use or application of products and/or materials referred to herein.
- (5) We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray. Combination sprays can induce crop phytotoxicity.
- (6) Apply at times of low light intensity, early morning or preferably during the evening. Avoid applications to wet leaves, stressed crops and on exceptionally hot, dry or sunny days. Overcast, calm and cool conditions are best.
- (7) E.&O.E.