



Suggested Nutrient Applications for:

Latest Update: 24/11/10



Wine Grapes

Growth Stage		Bud Burst	4Wk Post Bud Burst	6Wk Post Bud Burst	Pre-Flowering	Fruit Set	Pea Size Berries	Bunch Closure	Verasion	Post Harvest
Product	% Nutrient (N:P:K) w/v									
Firmrite SprayCal N	12.5 N, 18 Ca	30		30	30	Litres/Hectare				
Maxi Phos Injecta 23	16 N, 23 P		30							
Smartrace Zinc/Manganese	5 Zn, 5 Mn, 5.4 S		4 ←	5 ←	5 ←	5 ←				
Result Plus	14: 8: 10 + TE		4 ←	5 ←	5 ←	5 ←				
Boron 15	6 N, 15 B				2 ←	2 ←				
32.3	32 N, 0.3 HA	= * PRODUCT TO PRODUCT COMPATIBILITY								50#
Firmrite B	11Ca, 1.3B						5	5		

- * = Product to Product Compatibility when diluted with water at correct label rates
- On Merlot apply 2 applications of Moly Six (6% Mo, 0.4% B) prior to flowering @ 1lt/ha, is compatible with Smartrace Zinc/Manganese and Result Plus
- # Apply 32.3 3 weeks after Maxiphos Injecta 23 application

Minimum Water Rate: Foliar Application 1:50 Fertigation 1:100 = Foliar (Lt/Ha) = Fertigation (Lt/Ha)

Important Notes:

- (1) The product label should be consulted before use of any of the products referred to on this sheet, always refer to the label when applying SprayGro Liquid Fertilizer products.
- (2) To ensure correct nutrients are delivered to the plant, tissue testing should be carried out prior applying any nutrients to determine your crop situation. Crop damage is possible if trace elements are applied in excess more so than from deficiency.
- (3) This information sheet is a guide only as SprayGro Liquid Fertilizers neither makes, nor authorizes any agent or representative to make any other warranties, express or implied, other than those stated on the product label.
- (4) SprayGro Liquid Fertilizers shall not be liable for any results, loss, or damage whatsoever, whether consequential or otherwise through the use or application of products and/or materials referred to herein.
- (5) We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray. Combination sprays can induce crop phytotoxicity.
- (6) Apply at times of low light intensity, early morning or preferably during the evening. Avoid applications to wet leaves, stressed crops and on exceptionally hot, dry or sunny days. Overcast, calm and cool conditions are best.
- (7) E.&O.E.