

CAUTION

The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
- Use the correct label spray rates of any pesticide used with Spraygro product.
- **DO NOT** use when crop is under heavy stress from pests or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **DO NOT** spray to visible droplet coalescence. **DO NOT** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is imminent.

Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

12 SR-2



SMARTTRACE[®]

CHELATED ELEMENTS

SPRAYTRACE[®] 4

GUARANTEED MINIMUM ANALYSIS

Magnesium.....2% w/v Zinc.....2% w/v
 Iron.....2% w/v Manganese.....2% w/v

Trace elements as lignosulfonate chelates

20 Litre NET 200 Litre NET 1000 Litre NET



Application recommendations: Spraygro Smarttrace Spraytrace 4 is suitable for the following crops when trace element deficiency exists. Apply with sufficient water for thorough coverage of leaves. The rates are to be adjusted according to the tissue and/or soil analysis results.

BROADACRE: Rate of application per min water/hectare: 50 litres
 LUCERNE @ 3-5 litres/hectare. Apply prior to irrigation cycle immediately after first cut.
 COTTON @ 3-5 litres/hectare. All purpose foliar apply prior to irrigation.
 CEREALS @ 3-5 litres/hectare. Apply at 4 leaf stage and at tillering if required.

HORTICULTURE: Rate of application per min water/hectare: 300 litres
 CROPS @ 2-5 litres/hectare. Apply as needed or from when plants are 10-15cm in height. Apply every 14-21 days.
 CITRUS @ 5-6 litres/hectare. Apply 2 applications at bud burst 7-14 days apart. Apply at fruit set and filling.
 VINES @ 4-5 litres/hectare. Apply at regular intervals from bud burst.

ANALYSIS	w/v%
Magnesium as sulfate/lignosulfonate	2.00
Zinc (Zn) as lignosulfonate	2.00
Manganese (Mn) as lignosulfonate	2.00
Iron (Fe) as lignosulfonate	2.00
Sulfur as lignosulfonate/sulfate	6.2

Specific Gravity 1.25

WARNING: DO NOT apply to the foliage or fruit of sensitive crops such as apricots, nectarines, plums & pome fruit as phytotoxicity may occur.

Impurities
 Mo < 0.001%
 Se < 0.001%
 Cd < 1mg/kg
 Pb < 2mg/kg
 Hg < 0.2mg/kg

COMPATIBILITY: Highly incompatible with Amine salts of 2,4D or MCPA (excludes LVE versions). May block filters when combined with propiconazole. It is recommended to apply non-ionic wetter (e.g., BS1000) to improve compatibility. Add propiconazole first, then wetter and then **under agitation** add this product.

Liquid Fertilizer: Do not swallow, may be harmful. The spray from this product may act as an irritant. Avoid Inhalation and contact with eyes and skin.

WARNING
Hazards: May cause long-lasting harmful effects to aquatic life
Prevention: Wear protective gloves/protective clothing/eye protection/face protection. Avoid release to the environment.
Storage & Disposal: Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.



SPRAYGRO LIQUID FERTILIZERS
 76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494
 PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
 PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au

PRODUCED AND MANUFACTURED IN AUSTRALIA

NOT APPLICABLE 1000 LITRE CONTAINERS



DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect to any loss or damage including consequential loss or damage which may be suffered or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon any descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro or upon customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or conditions concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.

Batch No. 9 333056 001372