

CAUTION

The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
- Use the correct label spray rates of any pesticide used with Spraygro product.
- **DO NOT** use when crop is under heavy stress from pests or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **DO NOT** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is imminent.

Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

14 BS-4

LIQUID FERTILIZER

FIRMRITE™ CB

HIGH CALCIUM SOLUTION DESIGNED TO INCREASE
FRUIT DEVELOPMENT, FIRMINESS & MATURITY. CALCIUM
STRENGTHENS CELL WALLS IN FRUIT & VEGETABLES

16% w/v CALCIUM 0.2% w/v BORON

20 Litre NET 200 Litre NET 1000 Litre NET

HORTICULTURE: Foliar applications per min water/hectare generally required: 300L. Tree crops may require higher rates (500-1000L).

CROP	FOLIAR RATES L/Ha	COMMENTS
ASPARAGUS	2-4L	Apply 7 days from 1st fern, Repeat at spear emergence.
AVOCADOS	3-7L	Apply 3 weeks prior to flowering. Repeat every 3 weeks from fruit set to harvest.
BROCCOLI	2-4L	Apply every 2 weeks until head formation.
CAPSICUM	4-5L	Apply every 2 weeks until flowering. Repeat at regular intervals during fruiting.
CELERY	2-4L	Apply every 2 weeks. Stop addition 1 week before harvest.
CITRUS	5-6L	Apply prior to bloom, at petal fall and later as required.
COTTON	5-6L	Apply prior to 1st flowering. Repeat at boll formation
CUCUMBERS & CUCURBITS	2-5L	Apply prior to flowering with further applications at regular intervals. Do not apply a week before harvest.
GRAPE VINES	4-6L	Apply prior to flowering with regular applications until 2-3 weeks before harvesting. Minimum water 500L/ha.
LETTUCE	5-10L	Apply from emergence, continue 14 day intervals until harvest.
NUT CROPS	2-4L	Apply 2 weeks post bloom, repeat at regular intervals during nut development. Repeat post-harvest.

CROP	FOLIAR RATES L/Ha	COMMENTS
OLIVES	3-4L	Apply 2 weeks prior to fruit set, at fruit set and further applications as required.
POME FRUIT	5-10L	Apply at petal fall. Continue every 14 days to harvest.
POTATOES	4-5L	Apply 3 weeks after emergence. Repeat at hooking, canopy closure and bulking.
STONE FRUITS	5-10L	Apply post-bloom. Continue from fruit set at regular intervals.
STRAWBERRIES	2-4L	Apply 2 applications prior to flowering with further applications at regular intervals.
TOMATOES	7-16L	Apply at flowering then at 14 day intervals.
TROPICAL FRUIT	2-4L	Spray before bloom. Repeat at fruit set until 7 days before harvest.
VEGETABLES	2-5L	Apply 3 weeks after emergence or post-transplant.
ZUCCHINI	4-5L	Two applications prior to flowering. Repeat during fruit elongation until 1 week before harvest.

ANALYSIS	w/v%
Calcium (Ca) as calcium chloride	16
Boron (B) as borate	0.2

Specific Gravity 1.34 ± 0.01

Impurities
 Mo < 0.001%
 Se < 0.001%
 Cd < 1mg/kg
 Pb < 1mg/kg
 Hg < 0.2mg/kg

Liquid Fertilizer: Do not swallow, may be harmful. The spray from this product may act as an irritant. Avoid Inhalation and contact with eyes and skin.

WARNING
Hazards: Causes serious eye irritation.
Prevention: Wash hands thoroughly after handling. Wear protective gloves/protective clothing/eye protection/face protection.
Response: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do – continue rinsing. If eye irritation persists get medical advice/attention.
Storage & Disposal: Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.



NOT APPLICABLE 1000 LITRE CONTAINERS

SPRAYGRO LIQUID FERTILIZERS
 76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494
 PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
 PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au
PRODUCED AND MANUFACTURED IN AUSTRALIA

Batch No. 9 333056 002638

DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and/or user of Spraygro products ("customer"), in respect to any loss or damage including consequential loss or damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon any descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. 4) the customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.