

CAUTION

The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
- Use the correct label spray rates of any pesticide used with Spraygro product.
- **DO NOT** use when crop is under heavy stress from pests or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **DO NOT** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is imminent.

Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer"), in respect to any loss or damage including consequential loss or damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon any descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. 4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorized officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.



LIQUID FERTILIZER

FIRMRITTE™ CMB

A SOLUTION OF MICRO AND MACRO TRACE ELEMENTS
DESIGNED TO INCREASE THE YIELD OF CROPS BY
PROVIDING BALANCED CROP NUTRITION AND VIGOUR

10.5% NITROGEN, 21% CALCIUM,
3% MAGNESIUM, 0.1% BORON w/v

20 Litre NET 200 Litre NET 1000 Litre NET

CROP	FOLIAR RATES L/Ha	FERTIGATION RATES L/Ha	COMMENTS
NUT CROPS	2-4L	7-10L	Apply 2 weeks post bloom, repeat at regular intervals during nut development. Repeat post-harvest.
OLIVES	3-4L	8-10L	Apply 2 weeks prior to fruit set, at fruit set and further applications as required.
POME FRUIT	2-4L	7-10L	Apply at petal fall. Continue every 14 days to harvest.
POTATOES	4-5L	9-12L	Apply 3 weeks after emergence. Repeat at hooking, canopy closure and bulking.
STONE FRUITS	2-4L	7-20L	Apply post-bloom. Continue from fruit set at regular intervals.
STRAWBERRIES	2-4L	7-10L	Apply 2 applications prior to flowering with further applications at regular intervals.
TOMATOES	4-5L	9-12L	Apply every 14-21 days from 2nd truss to harvest.
TROPICAL FRUIT	2-4L	7-10L	Spray before bloom. Repeat at fruit set until 7 days before harvest.
VEGETABLES	2-5L	7-10L	Apply 3 weeks after emergence or post-transplant. Repeat at regular intervals.
ZUCCHINI	4-5L	9-12L	Two applications prior to flowering. Repeat during fruit elongation until 1 week before harvest

ANALYSIS	w/v%
Calcium (Ca) as calcium nitrate	10.7
Calcium (Ca) as calcium chloride	10.3
Calcium (Ca) as total	21.0
Magnesium (Mg) as magnesium nitrate	3.0
Boron (B) as borate	0.1

Specific Gravity 1.58 ± 0.01

Impurities
Mo < 0.001%
Se < 0.001%
Cd < 1mg/kg
Pb < 2mg/kg
Hg < 0.2mg/kg



HORTICULTURE: Foliar applications per min water/hectare generally required: 150-300L. Tree crops may require higher rates (500-1000L).

CROP	FOLIAR RATES L/Ha	FERTIGATION RATES L/Ha	COMMENTS
ALMONDS	2-4L	7-10L	Apply 2 weeks post-bloom. Repeat 2-3 times every 4 weeks.
APPLES	5-7L	10-20L	Apply at petal fall. Continue at 7-14 day intervals until harvest.
ASPARAGUS	2-4L	7-10L	Apply 7 days from 1st fern. Repeat at spear emergence.
AVOCADOS	2-4L	7-10L	Apply 3 weeks prior to flowering. Repeat every 3 weeks from fruit set to harvest.
BANANAS	3-4L	8-10L	Preflowering apply up to 3-4L/ha, after flowering apply only 2L/ha via foliar. Repeat if necessary. (Caution: surface of recently emerged fruit is particularly prone to marking damage from spray applications made to very rapidly growing fruit in hot conditions with direct light. Recommended to bag all emerged bunches prior to spraying to minimise risk of fruit marking).
BROCCOLI	2-4L	7-10L	Apply every 2 weeks until head formation.
CAPSICUM	4-5L	9-12L	Apply every 2 weeks until flowering. Repeat at regular intervals during fruiting.
CELERY	2-4L	7-10L	Apply every 2 weeks. Stop application 1 week before harvest.
CITRUS	5-8L	10-20L	Apply prior to bloom, at petal fall and later as required.
COTTON	5-6L	10-20L	Apply prior to 1st flowering. Repeat at boll formation.
CUCUMBERS & CUCURBITS	2-5L	8-12L	Apply prior to flowering with further applications at regular intervals do not apply a week before harvest.
GRAPE VINES	4-6L	9-15L	Apply prior to flowering with regular applications until 2-3 weeks before harvesting.
LETTUCE	3-4L	8-10L	Apply from emergence, continue 14 day intervals until harvest.

Liquid Fertilizer: Do not swallow, may be harmful. The spray from this product may act as an irritant. Avoid Inhalation and contact with eyes and skin.

WARNING
Hazards: Causes serious eye irritation.
Prevention: Wash hands thoroughly after handling. Wear protective gloves/protective clothing/eye protection/face protection.
Response: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do – continue rinsing. If eye irritation persists get medical advice/attention.
Storage & Disposal: Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.



SPRAYGRO LIQUID FERTILIZERS
76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494
PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au
PRODUCED AND MANUFACTURED IN AUSTRALIA

NOT APPLICABLE 1000 LITRE CONTAINERS