

CAUTION

The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
- Use the correct label spray rates of any pesticide used with Spraygro product.
- **DO NOT** use when crop is under heavy stress from pests or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **DO NOT** spray to visible droplet coalescence. **DO NOT** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is imminent.

Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer"), in respect to any loss or damage including consequential loss or damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon any descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. 4) the customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.



LIQUID FERTILIZER

FIRMRITE™ CP

High analysis Phosphorus solution with Calcium to promote plant development, hasten maturity and increase firmness

N P K
0 : 14 : 0 %w/v
+4% Calcium

20 Litre NET 200 Litre NET 1000 Litre NET

- Foliar rate of application per min water/hectare:** 300 litres
- ASPARAGUS @ 3-5 litres/hectare applied to ferns prior to senescence x3 applications 10-14 days apart.
 - CUCURBITS @ 5 litres/hectare x3-5 applications at 7 day intervals commencing at first flowers through fruit expansion.
 - LETTUCE @ 5 litres/hectare commencing 10-14 days after transplanting or emergence with 7-14 days between applications.
 - ONIONS @ 5 litres/hectare during bulb tillering x1-2 applications.
 - PINEAPPLE @ 5 litres/hectare commencing at early fruit development.
 - POTATOES @ 10 litres/hectare at tuber initiation to increase tuber numbers and 2-3 applications to promote tuber size during bulking.
 - STRAWBERRIES @ 5 litres/hectare from start of flowering at 7-10 day intervals
- Foliar rate of application to following crops per min water/hectare:** 500-1000 litres
- APPLES/PEARS @ 12 litres/hectare x4-5 applications at 7-10 day intervals commencing 1 week after petal fall. Colour enhancement via applying 10-15 litres/ha, red colour in apples and stone fruit is greater, Apply when fruit has started to change colour. Normally 2-3 weeks before harvest.
 - BRASSICAS @ 10 litres/hectare from stem extension/head development at 7-10 day intervals.

- CITRUS @ 10 litres/hectare at fruit set with 1-2 applications at 7 day intervals.
- KIWI FRUIT @ 10 litres/hectare commencing post blossom x4 applications.
- PAW PAW @ 5 litres/hectare commencing at early fruit development x3 applications.
- PERSIMMONS @ 10 litres/hectare after flowering and repeat 10-14 days later.
- STONE FRUIT @ 10 litres/hectare apply 3-7 days after petal fall x3 applications applied at 7-10 day intervals.
- FIELD TOMATOES @ 5 litres/hectare commencing at flowering on second truss onwards x2-4 applications at 10-14 day intervals.
- VINES/GRAPES @ 10 litres/hectare from fruit set at 7-10 day intervals x3-5 applications.

ANALYSIS	w/v%
Calcium (Ca) as calcium dihydrogen phosphate	4.9
Phosphorus (P) as calcium dihydrogen phosphate	14.0

Specific Gravity 1.29 ± 0.01

Impurities
Mo < 0.001%
Se < 0.001%
Cd < 1mg/kg
Pb < 1mg/kg
Hg < 0.2mg/kg
F < 40g/kg P

WARNING: Do not mix with other Ag chemicals when used for colour enhancement in crops.

Liquid Fertilizer: Do not swallow, may be harmful. The spray from this product may act as an irritant. Avoid Inhalation and contact with eyes and skin.

DANGER
Hazards: Causes skin irritation. Causes serious eye damage.
Prevention: Wash hands thoroughly after handling. Wear protective gloves/protective clothing/eye protection/face protection.
Response: Immediately call a POISON CENTER or doctor. Take off contaminated clothing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do – continue rinsing. If skin irritation occurs: Get medical advice/attention.
Storage & Disposal: Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.



SPRAYGRO LIQUID FERTILIZERS
76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494
PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au

PRODUCED AND MANUFACTURED IN AUSTRALIA

NOT APPLICABLE 1000 LITRE CONTAINERS