

CAUTION

LIQUID FERTILIZER

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FOUR 7 TRACE

GUARANTEED MINIMUM ANALYSIS

| | | | |
|-------------|-----------|---------------|-----------|
| SULFUR..... | 7.0% w/v. | NITROGEN..... | 7.0% w/v. |
| IRON..... | 7.0% w/v. | ZINC..... | 7.0% w/v. |
| COPPER..... | 0.5% w/v | | |

20 Litre NET 200 Litre NET 1000 Litre NET

FOUR 7 TRACE suitable for the following crops where a Zinc and Iron deficiency occurs. The rates must be adjusted according to tissue and or soil analysis results.

BROADACRE: Rate of application per min water/hectare: 50 litres.

- Cereals @ 1-2 litres/hectare
- Cotton & Soybeans @ 1-2 litres/hectare or irrigated crops apply just prior to irrigation to prevent crop shed and increase yield.

COMPATIBILITY: Highly incompatible with Amine salts such as 2,4D and MCPA (excludes LVE). May block filters when combined with propiconazole. It is recommended to apply non-ionic wetter (e.g., BS1000) to improve compatibility. Add propiconazole first, then wetter and then under agitation add this product.

CROP SAFETY: Avoid spraying sulfate trace elements when conditions cause excessive evaporation and leaf burn. Be aware that additional nitrogen fertilizer to this product may increase the risk of phytotoxicity.

WARNING: DO NOT apply to the foliage or fruit of sensitive crops such as apricots, nectarines, plums & pome fruit as phytotoxicity may occur.

| ANALYSIS | w/w% |
|-------------------------------|------|
| Sulfur (S) as sulfate | 7.0 |
| Nitrogen (N) as urea | 7.0 |
| Iron (Fe) as iron sulfate | 7.0 |
| Zinc (Zn) as zinc sulfate | 7.0 |
| Copper (Cu) as copper sulfate | 0.5 |

Specific Gravity 1.37 ± 0.01

Impurities
Mo < 0.001%
Se < 0.001%
Cd < 1mg/kg
Pb < 2mg/kg
Hg < 0.2mg/kg
biuret < 10g/kg N

DANGER
Hazards: Harmful if swallowed. Harmful in contact with skin. Causes skin irritaiton. Causes serious eye damage. Toxic to aquatic life. Harmful to aquatic life with long-lasting effects.
Prevention: Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Avoid release to the environment. Wear protective gloves/protective clothing/eye protection/face protection.
Response: Immediately call a POISON CENTER or doctor. Rinse mouth. Take off contaminated clothing. Wash contaminated clothing before reuse and flush skin and hair with running water. IF SWALLOWED: Immediately call a POISON CENTER or doctor. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do – continue rinsing. For advice, contact a Poisons Information Centre (e.g. phone Australia 13 11 26; New Zealand 0800 764 766) or a doctor (at once). Collect spillage.
Storage & Disposal: Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for information.



NOT APPLICABLE 1000 LITRE CONTAINERS

SPRAYGRO LIQUID FERTILIZERS
 76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494
 PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
 PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au

PRODUCED AND MANUFACTURED IN AUSTRALIA



DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect to any loss or damage including consequential loss or damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon any descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. 4) the customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.

Batch No. 333056 001693 9

The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
- Use the correct label spray rates of any pesticide used with Spraygro product.
- DO NOT use when crop is under heavy stress from pests or inadequate moisture levels.
- DO NOT apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- DO NOT spray to run off. DO NOT spray to visible droplet coalescence. DO NOT allow concentrated spray mist to run on fruit or leaves.
- DO NOT apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is imminent.

Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.