

26 BS-6

CAUTION



LIQUID FERTILIZER

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

BOLLY 5:3

TO BE USED WHERE A BORON-MOLYBDENUM
DEFICIENCY EXISTS

N P K
3 0 0 + 5 B, 3 Mo w/v%

20 Litre NET 200 Litre NET 1000 Litre NET

Bolly 5:3 application recommendations: Apply to the following crops where a Boron-Moly deficiency exists. Minimum water rate is 1:50 for foliar application and 1:100 for fertigation.

CROP	FOLIAR RATES L/Ha	FERTIGATION RATES L/Ha	COMMENTS
ALMONDS	2-4L	9-15L	For best results apply onto new foliage when symptoms are exposed.
APPLES	2-3L	6-12L	For best results apply onto new foliage when symptoms are exposed.
ASPARAGUS	1-2L	3-6L	Apply as symptoms are exposed.
AVOCADOS	3-6L	9-15L	Apply as symptoms are exposed or apply at 4-5 leaf stage.
BANANAS	1.5-3L	6-9L	Apply as symptoms are exposed.
BROCCOLI	1-2L	3-6L	Apply as symptoms are exposed or apply at 4-5 leaf stage.
CAPSICUM	1-2L	3-6L	Apply as symptoms are exposed or apply at 4-5 leaf stage.
CELERY	0.5-3L	3-9L	Apply as symptoms are exposed or apply at 4-5 leaf stage.
CEREALS	2L	9-15L	Apply as symptoms are exposed.
CITRUS	3-5L	9-15L	For best results apply onto new foliage when symptoms are exposed.
COTTON	3-6L	9-15L	For best results apply onto new foliage when symptoms are exposed.
GRAPE VINES	3-5L	9-13L	For best results apply onto new foliage when symptoms are exposed.
LEGUMES	2-4L	9-15L	Apply when early growth is approx 150mm tall to ensure effective movement into nitrogen fixing nodules.
MELONS	1-2L	3-6L	Apply as symptoms are exposed or apply at 4-5 leaf stage.
NUT CROPS	2-4L	9-15L	For best results apply onto new foliage when symptoms are exposed.

CROP	FOLIAR RATES L/Ha	FERTIGATION RATES L/Ha	COMMENTS
OLIVES	2-4L	9-15L	For best results apply onto new foliage when symptoms are exposed.
LUCERNE	1.2-2L	1.2-2L	Apply when early growth is approx 150mm tall to ensure effective movement into nitrogen fixing nodules. Apply at least 2 weeks before cutting. Minimum water dilution 1:200
PASTURE	0.8-1.2L	1.6-2.4L	
POPPIES	2-4L	9-15L	Apply from row cover till capule development as needed.
POTATOES	2-4L	9-15L	Apply as symptoms are exposed or apply at 4-5 leaf stage.
STONE FRUIT	0.5-2L	3-6L	For best results apply onto new foliage when symptoms are exposed.
STRAWBERRIES	1-2L	3-6L	Apply as symptoms are exposed or apply at 4-5 leaf stage.
TOMATOES	2-4L	9-15L	Apply as symptoms are exposed or apply at 4-5 leaf stage.
TREE CROPS	2-4L	9-15L	For best results apply onto new foliage when symptoms are exposed.
VEGETABLES	1-2L	3-6L	Apply as symptoms are exposed or apply at 4-5 leaf stage.
ZUCCHINI	1-2L	3-6L	Apply as symptoms are exposed or apply at 4-5 leaf stage.

ANALYSIS	w/v%
NITROGEN N	3.0
Boron B	5.0
Molybdenum Mo	3.0

Specific Gravity 1.22

Impurities
Se < 0.001%
Cd < 1mg/kg
Pb < 3mg/kg
Hg < 0.2mg/kg

The rates must be adjusted according to the tissue and/or soil analysis results.

WARNING: Excessive use of molybdenum (Mo) can be harmful to stock. Plant levels of Mo can be high for 4 weeks after application. It is Advisable to keep stock off treated areas for this period. Molybdenum application may induce copper deficiency in grazing animals. This is most likely to occur on sandy soils low in copper. Check rate and frequency of molybdenum use with appropriate authorities.

Liquid Fertilizer: Do not swallow (schedule 5 poison). The spray from this product may act as an irritant. Avoid Inhalation and contact with eyes and skin.

This product is not classified as Hazardous according to the criteria of Safe Work Australia. Storage & Disposal: Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.



DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect to any loss or damage including consequential loss or damage which may be suffered or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon any descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. 4) the customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. (6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.

Batch No. 333056 002515 9

The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
- Use the correct label spray rates of any pesticide used with Spraygro product.
- DO NOT use when crop is under heavy stress from pests or inadequate moisture levels.
- DO NOT apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- DO NOT spray to run off. DO NOT spray to visible droplet coalescence. DO NOT allow concentrated spray mist to run on fruit or leaves.
- DO NOT apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is imminent.

Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.