

## CAUTION

The following conditions must be observed in order to apply SprayGro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use SprayGro product under conditions of optimum plant growth including highest humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day.
- When mixing with other chemicals, always mix a small quantity (e.g. bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiables or flowables to water in the mix and wet, dissolve or disperse before adding SprayGro product.
- Use the correct label spray rates of any pesticide used with SprayGro product.
- **DO NOT** use when crop is under heavy stress from pests, heat or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **Do not** spray to visible droplet coalescence. **Do not** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day.

**NOTE:** We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

**DISCLAIMER/CONDITIONS OF SALE:** 1) Foliar Fertilizers Pty Ltd trading as SprayGro Liquid Fertilizer and its servants and agents ("SprayGro") shall be under no liability to the purchaser and or user of SprayGro products ("customer") in respect of any loss or damage including consequential loss or damage which may be suffered, incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the SprayGro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by SprayGro or upon and descriptions, illustrations or specifications contained in SprayGro's Product Profile or any other document produced by SprayGro. 3) The customer understands that any representation of the type set out above is on the basis that the customer must independently verify the accuracy thereof. 4) The customer shall at all times indemnify and hold harmless SprayGro from and against all loss (including reasonable legal costs and expenses) or liability reasonably incurred or suffered by SprayGro arising from any proceedings or threatened proceedings against SprayGro by or involving the customer. 5) SprayGro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of SprayGro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from SprayGro or our Distributor.

**BATCH No.**



9 1335056 00329 1



## LIQUID FERTILIZER

# 32-3

N P K

32 : 0 : 0 + Humic Acid

TOTAL NITROGEN (N) ..... 32.0% W.V.  
 NITROGEN AS UREA ..... 16.0% W.V.  
 NITROGEN AS AMMONIUM ..... 8.0% W.V.  
 NITROGEN AS NITRATE ..... 8.0% W.V.  
 HUMIC ACID ..... 0.3% W.V.

32-3 IS NOT RECOMMENDED AS A FOLIAR SPRAY IN HORTICULTURE

20 Litre NET  200 Litre NET  1000 Litre NET

### BROADACRE:

**Foliar application rate: in minimum 50-100 litres water/hectare.**

FIELD CROPS @ Apply at 10-40 litres per hectare from 4 leaf stage for improved yield. Apply at early Boot Stage to increase Protein.

COTTON @ Apply at 2-5 litres per hectare from 4-6 leaf stage. Apply at 10-20 litres per hectare at boll development.

### HORTICULTURE:

**DO NOT USE AS A FOLIAR SPRAY IN HORTICULTURE.**

APPLICATION RATE: Water rate 1000 litres/hectare.

FERTIGATION @ 20-30 litres per hectare. Inject into irrigation systems including drip, trickle, flood. Do not wet the foliage of the plant. Suited to a wide variety of crops including vegetables, trees, tomatoes, citrus, bananas and apples.

**WARNING:** Repeated foliar applications of this fertilizer may cause damage to citrus, avocados and pineapples (due to biuret).

**CAUTION:** 32.3 is NOT compatible with 2,4D Amine and /or MCPA salts. Use LVEs instead.

GRAPES @ Fertigation, 10-20 litres per hectare. Inject into the water system during the second half of the irrigation cycle.

TURF @ 500-700 mls per 100m<sup>2</sup> water dilution 1:100.

**NOTE:** When foliar applying Nitrogen to crops, the lower the rate of water applied will improve results, **however**, crops under any moisture stress, or under high temperatures or windy situations will be open to potential fertilizer (salt) burn. Use lower rates of water at your own discretion.

ANALYSIS	AUSTRALIA W/V %	INTERNATIONAL W/W %
Nitrogen N	32	26.44
Humic Acid	0.3	0.25

Specific gravity 1.21 pH 8.5-9.5

### STORAGE & DISPOSAL:

Do not contaminate water, food or feed by storage or disposal. Triple rinse container, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

### SPRAYGRO LIQUID FERTILIZERS

76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • P.O. BOX 494 PORT ADELAIDE BUSINESS CENTRE  
 PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496  
 PHONE: (08) 8447 7266 • FAX: (08) 8240 1844 • Email: spraygro@spraygro.com.au • www.spraygro.com.au

**PRODUCED AND MANUFACTURED IN AUSTRALIA**



NOT APPLICABLE TO 1000 LITRE CONTAINERS