


40 HN-4



Pick 15-42

BROADACRE: Barley, Canola, Cotton, Legumes, Maize, Oats, Rice, Sorghum, Triticale, Wheat & Pasture Crops.

Foliar Rate/Ha	Fertigation/Ha	Comments
3-6L	N/A	Apply in a minimum of 300-600L final spray volume. Apply from mid tillering or late cabbage stage or as required. For cotton: Apply from first flowering until 2 weeks before harvest. Assists in delaying leaf senescence.

NUT TREES: Almond, Pistachio & Walnut.

2 - 5L	10-15 L	Foliar: Apply in a minimum of 400-1000L final spray volume. Apply with compatible crop protection sprays. Fertigation as required. Best applied via fertigation or post-harvest to increase potassium levels.
--------	---------	---

FRUIT TREES: Apple, Cherry, Nectarine, Peach & Pear.

N/A	10-15 L	DO NOT apply as a foliar on stone fruit. Fertigation as required. Best applied via fertigation or post-harvest to increase potassium levels.
-----	---------	--

EVERGREEN TREE CROPS: Avocado, Citrus, Macadamia & Lychee.

2 - 4L	10-20 L	Foliar Apply in a minimum of 300-600L final spray volume. Apply to recently hardened spring flush or during active growing period, fruit fill and ripening to accelerate fruit maturity.
--------	---------	--

FRUITING VEGETABLES: Capsicum, Cucurbits, Eggplant, Tomatoes (field), Watermelon & Pumpkins.

3-5L	15-30 L	Foliar: Apply in a minimum of 300-500L final spray volume. 1st application prior to flowering repeated at 14 day intervals during filling and ripening.
------	---------	---

LEAFY VEGETABLES: Endive, Fennel Lettuce, Broccoli, Cabbage, Cauliflower, Kale & Herbs.

Foliar Rate/Ha	Fertigation/Ha	Comments
3-5L	5-10 L	Foliar: Apply in a minimum of 300-500L final spray volume. Foliar spray 10-14 days post-transplant. Repeat at 7-14 day intervals, dependent on severity of deficiency.

ROOT VEGETABLES: Beetroot, Carrot, Leek, Onion, Potato, Radish & Sweet Potato.

3-5L	15-30L	Foliar: Apply in a minimum of 450-750L final spray volume. Apply as required to supply potassium to aid maturity. Apply 2 sprays 14 days apart on onions during bulb formation. For potatoes: Apply at planting and then from hooking until harvest.
------	--------	--

VINE and BERRY CROPS: Blueberry, Strawberry, Raspberry, Wine & Table grapes.

2 - 4L	15-20L	Foliar: Apply in a minimum of 300-400L final spray volume. Apply as required. DO NOT exceed 1x concentration of per hectare rate. Care should be taken when applying to grapes. When applied under certain conditions, K Blast Neutral may disturb the bloom - always test spray a small area and observe prior to total area application.
--------	--------	--

ANALYSIS	w/v%
Potassium (K) as potassium phosphate	15
Phosphorus (P) as potassium phosphate	42

Specific Gravity 1.62 ± 0.01
pH at 10% dilution 10.5 - 11.5



Liquid Fertilizer: Do not swallow, may be harmful. The spray from this product may act as an irritant. Avoid Inhalation and contact with eyes and skin.

Not classified as a hazardous substance according to the criteria of Safe Work Australia.

Prevention: Wear protective gloves/protective clothing/eye protection/face protection.

Storage & Disposal: Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

SPRAYGRO LIQUID FERTILIZERS
76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494
PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au

PRODUCED AND MANUFACTURED IN AUSTRALIA

Impurities

Mo < 0.001%
Se < 0.001%
Cd < 1mg/kg
Pb < 1mg/kg
Hg < 0.2mg/kg
F < 40g/kg P

Batch No.



DISCLAIMER/CONDITIONS OF SALE: (1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect to any loss or damage including consequential loss or damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. (2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon any descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. (3) The customer understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. (4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. (5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. (6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.

The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.
• Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
• Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
• Use the correct label spray rates of any pesticide used with Spraygro product.
• DO NOT use when crop is under heavy stress from pests or inadequate moisture levels.
• DO NOT apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
• DO NOT spray to run off. DO NOT spray to visible droplet coalescence. DO NOT allow concentrated spray mist to run on fruit or leaves.
• DO NOT apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is imminent.
Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.