

CAUTION

The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
- Use the correct label spray rates of any pesticide used with Spraygro product.
- **DO NOT** use when crop is under heavy stress from pests or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **DO NOT** spray to visible droplet coalescence. **DO NOT** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is imminent.

Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and/or user of Spraygro products ("customer"), in respect to any loss or damage including consequential loss or damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon any descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. 4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.



44 NH-5



Application recommendation for Result Plus 14-8-10. Apply with sufficient water for thorough coverage of leaves when foliar applying.

HORTICULTURE: Foliar applications per min water/hectare generally required: 150-350L. Tree crops may require higher rates (500-1000L)

CROP	FOLIAR RATES L/Ha	FERTIGATION RATES L/Ha	COMMENTS
AVOCADOS & TROPICAL FRUIT	4L	15L	Apply at 7-14 day intervals from fruit set through to harvest.
ALMONDS	2-3L	15-30L	Apply post-bud burst at 14 day intervals until hull split.
ASPARAGUS	2-4L	7-10L	Apply at spear emergence until harvest.
CITRUS	2-4L	15-30L	Apply post-flowering until fruit expansion.
CAPSICUM	3-4L	10-30L	Apply post-planting until mid fruiting.
BRASSICA'S	4L	10-15L	Apply at early head development to harvest every 14 days.
CUCURBITS	2-4L	15-20L	Apply at fruit set onwards every 7-14 days until harvest.
GARLIC AND ONIONS	3-4L	12-15L	Apply from when sufficient leaf area exists to intercept spray and at bulb development through to harvest.
GRAPES	3-5L	15-30L	Apply post-bud burst until veraison.
LEAFY GREENS	2-3L	8-12L	Apply every 7-14 days from 4 leaf stage until harvest.
OLIVE TREES	3-4L	15-20L	Apply from early through to late fruiting.
PAW PAW	2-3L	8-12L	Apply at end of flowering and 21 days after flowering as required.
PEANUTS	n/a	10-20L	Apply at planting.

CROP	FOLIAR RATES L/Ha	FERTIGATION RATES L/Ha	COMMENTS
PINEAPPLES	2-3L	5-10L	Apply at 14-21 day intervals from fruit set through to harvest.
POTATOES	4-5L	12-20L	Apply every 7-14 days from tuber initiation to harvest.
STRAWBERRIES	2-4L	10-20L	Apply post-planting at regular intervals.
TOMATOES	2-3L	10-15L	Apply at early to late vegetative growth.
VEGETABLES	4-6L	10-15L	Apply at regular intervals during growing season.

BROADACRE: Foliar rate of application per min water/hectare: 50 litres.
 COTTON @ 6-9 litres/hectare at 4-6 leaf stage and during mid to late flowering.
 CEREALS @ 6-9 litres/hectare when sufficient leaf area exists to spray.
 LEGUME GRAINS @ 4 litres/hectare during pod development.
 TURF @ 400-500ml/100m². Water dilution 1:100

ANALYSIS	w/v%	ANALYSIS	w/v%
Nitrogen (N) as urea	11.7	Iron (Fe) as EDTA	0.03
Nitrogen (N) as ammonia	2.3	Zinc (Zn) as EDTA	0.02
Nitrogen (N) as total	14	Manganese (Mn) as EDTA	0.03
Phosphorus (P) as phosphate	8	Copper (Cu) as EDTA	0.02
Potassium (K) as potassium phosphate	10	Boron (B) as Borate	0.017
		Molybdenum (Mo) as Molybdate	0.002
		Cobalt (Co) as EDTA	0.001
Specific Gravity 1.29 ± 0.01			



Liquid Fertilizer: Do not swallow, may be harmful. The spray from this product may act as an irritant. Avoid Inhalation and contact with eyes and skin.

Not classified as a hazardous substance according to the criteria of Safe Work Australia.
Prevention: Wear protective gloves/protective clothing/eye protection/face protection.
Storage & Disposal: Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

Impurities
 Se < 0.001%
 Cd < 1mg/kg
 Pb < 2mg/kg
 Hg < 0.2mg/kg
 F < 40g/kg P
 Biuret <8g/kg N

SPRAYGRO LIQUID FERTILIZERS
 76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494
 PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
 PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au

PRODUCED AND MANUFACTURED IN AUSTRALIA