



Batch No. 333056 9 003949  
 (9) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.  
 Disclaimer/Conditions of Sale: (1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect of any loss or damage including consequential loss or damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. (2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon any descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. (3) The customer understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. (4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. (5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product.

Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

- The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.
- Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
- Use the correct label spray rates of any pesticide used with Spraygro product.
- **DO NOT** use when crop is under heavy stress from pests or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **DO NOT** spray to visible droplet coalescence. **DO NOT** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is imminent.

## CAUTION

48 NH-4

**ZINTRON**  
**LIQUID FERTILIZER**  
**19-0-0 + 6.6 Zn**  
**CONTAINS Zn AS ZINC NITRATE FOR MAXIMUM FOLIAR UPTAKE.**

20 Litre NET  200 Litre NET  1000 Litre NET

Zintron applications recommendations: Apply when zinc deficiency exists. Horticulture - Foliar applications per minimum water/hectare spray to cover generally: 150-300L. Tree crops may require higher rates 500-1500L.

CROP	FOLIAR RATES L/Ha	COMMENTS
ALMONDS	1-3L	Apply at petal fall, repeat every 4 weeks, post harvest.
ASPARAGUS	1-4L	Apply at full fern. One application per year.
AVOCADO	1-3L	Best applied at Spring flush. Repeat as required.
BEANS	1-4L	Apply at sufficient foliage, repeat 10d up to full bloom.
BLUEBERRIES	0.5-1L	Apply from leaf expansion up to small fruits.
BROCCOLI	1-4L	Apply first at 150-200mm. Repeat 7-10 days.
BRUSSELS SPROUTS	1-2L	Apply first at 150-200mm. Repeat 7-10 days.
CABBAGE	1-4L	Apply at 10d to 2 weeks post thinning. Repeat if required.
CARROTS	1-4L	Apply at 100-150mm tall. Repeat in 10-14 days.
CELERY	1-2L	Apply at 75-150mm tall. Repeat in 10-14 days.
CITRUS	1-2L	Best applied at Spring and Autumn flush. Repeat if required.
CORN/MAIZE	1-2L	Apply when knee high and repeat before tasselling.
CUCUMBER	1-2L	Apply at first bloom and repeat 0.5L after each picking.
GARLIC	1-4L	Apply when 75-175mm tall and repeat as required.
GRAPES	1-3L	Apply from first leaves up to full bloom. Post harvest.
LETTUCE	1-2L	Apply at 10 days to 2 weeks post-thinning.
ONIONS	1-4L	Apply when 75-150mm tall. Repeat as required.
PEANUTS	1-2L	Apply 2-4 sprays from first bloom.
PISTACHIOS	1-2L	Apply at bud break to end of shoot elongation, dormant.
POME FRUIT	0.5-1L	Apply foliar post harvest, dormant only.
POTATOES	2-5L	Apply at hooking, canopy closure and tuber bulking.

The rates must be adjusted according to the tissue and/or soil analysis results.

CROP	FOLIAR RATES L/Ha	COMMENTS
STONE FRUIT	0.5-1L	Apply foliar post harvest, dormant only.
STRAWBERRIES	2-4L	Apply 2 weeks post planting and repeat 2-3 applications.
SUGARCANE	1-2L	Apply when sufficient foliage is present.
TOMATOES	1-4L	Apply from first truss and repeat as required.
WATERMELONS	0.5-1.5L	Apply up to first bloom.
ZUCCHINIS	1-3L	Apply 2 weeks post thinning and repeat 7-10 days.

**Broadacre - Foliar applications per minimum water/hectare 50L.**

CROP	FOLIAR RATES L/Ha	COMMENTS
CANOLA	1-3L	Apply from 4th leaf up to pod formation.
CEREALS	2-5L	Apply mid tillering to heading.
COTTON	1-2L	Apply 6-8th leaf stage and repeat at boll formation.
RICE	3-7L	Apply 35-55 days after emergence.

**FERTIGATION - May be applied at 7-20L/ha via overhead or dripper irrigation. Apply Zintron during middle 1/3 of irrigation cycle. Zintron can be applied with Firmrite Spray Cal N or UAN solutions.**

ANALYSIS	w/v%	Impurities
Zinc (Zn) as nitrate	6.6	Mo < 0.001%
Nitrogen (N) as nitrate	10.5	Se < 0.001%
Nitrogen (N) as ammonium	7.8	Cd < 1mg/kg
Nitrogen (N) as urea	0.7	Pb < 5mg/kg
Nitrogen (N) as total	19.0	Hg < 0.2mg/kg
		biuret < 0.36g/Kg N
		Specific Gravity 1.32 ± 0.01 pH 3-4



**WARNING**  
**Hazards:** Harmful if swallowed, Harmful in contact with skin. Causes skin and serious eye irritation. Toxic to aquatic life and harmful with long -lasting effects.  
**Prevention:** Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Avoid release to the environment. Wear protective gloves/protective clothing/eye protection/face protection.  
**Response:** Call a POISON CENTER or doctor if you feel unwell. Rinse mouth. Take off contaminated clothing. Wash contaminated clothing before use. IF SWALLOWED do NOT induce vomiting and Immediately call a POISON CENTER or doctor. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do – continue rinsing. If eye or skin irritation persists get medical advice/attention. For advice, contact a Poisons Information Centre (e.g. phone Australia 13 11 26; New Zealand 0800 764 766) or a doctor (at once).  
**Storage & Disposal:** Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.



NOT APPLICABLE 1000 LITRE CONTAINERS

**SPRAYGRO LIQUID FERTILIZERS**  
 76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494  
 PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496  
 PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au

**PRODUCED AND MANUFACTURED IN AUSTRALIA**