

50 GP-1

# CAUTION

The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
- Use the correct label spray rates of any pesticide used with Spraygro product.
- **DO NOT** use when crop is under heavy stress from pests or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **DO NOT** spray to visible droplet coalescence. **DO NOT** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is imminent.

Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

**DISCLAIMER/CONDITIONS OF SALE:** 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and/or user of Spraygro products ("customer"), in respect to any loss or damage including consequential loss or damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon any descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. 4) the customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.



# FULVIC ACID 12.5

## FOLIAR FERTILIZER BOOSTER

Suitable for use in conjunction with most liquid fertilizers in foliar applications as an uptake enhanced and complexing agent.

Analysis: 12.5% Fulvic acid

- 20 Litre NET
- 200 Litre NET
- 1000 Litre NET

ANALYSIS	w/v%
Fulvic acid	12.5
Potassium (K)	1.8
Total: Potassium Fulvate	14.3
Carbon (C)	8.0

Specific Gravity 1.08 ± 0.01 pH 5 to 6

**Impurities**  
 Mo < 0.001%  
 Se < 0.001%  
 Cd < 1mg/kg  
 Pb < 1mg/kg  
 Hg < 0.2mg/kg

### BENEFITS OF FULVIC ACID 12.5:

- Improves the uptake of nutrients by plants in foliar applications
- Promotes quicker germination and faster root and shoot growth.
- Promotes soil microbial activity.
- Complexing agent for micro-nutrients

### FOLIAR APPLICATION:

Apply 1-5L per hectare.

### MIXING WITH FERTILIZERS:

Fulvic acid 12-5 is compatible with most fertilizers when sufficient water is applied.  
 Recommended for blending with micronutrients such as the Bluestripe range of products.

### MIXING WITH PESTICIDES:

Fulvic acid 12-5 is compatible with most pesticides when sufficient water is applied.



Liquid Fertilizer: Do not swallow, may be harmful. The spray from this product may act as an irritant. Avoid Inhalation and contact with eyes and skin.

**Not classified as a hazardous substance according to the criteria of Safe Work Australia.**  
**Prevention:** Wear protective gloves/protective clothing/eye protection/face protection.  
**Storage & Disposal:** Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

SPRAYGRO LIQUID FERTILIZERS  
 76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494  
 PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496  
 PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au

PRODUCED AND MANUFACTURED IN AUSTRALIA

NOT APPLICABLE 1000 LITRE CONTAINERS