

# CAUTION

The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
- Use the correct label spray rates of any pesticide used with Spraygro product.
- **DO NOT** use when crop is under heavy stress from pests or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **DO NOT** spray to visible droplet coalescence. **DO NOT** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is imminent.

Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

**DISCLAIMER/CONDITIONS OF SALE:** (1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and/or user of Spraygro products ("customer") in respect to any loss or damage including consequential loss or damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. (2) The customer warrants that there has been no reliance upon any representation made by Spraygro or upon any descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. (3) The customer understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. (4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. (5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. (6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.





# Gripper™

e d t a c h e l a t e

# IRON w/v

6%

20 Litre NET  200 Litre NET  1000 Litre NET

**Application recommendations:** Gripper Iron is suitable for the following crops where Iron deficiency exists. Apply with sufficient water for thorough leaf coverage.

CROP		RATE/ha	Minimum dilution	Comments
ALMONDS	Foliar	2-3L	1:100	Apply as required to maintain crop colour and vigour. Repeat early fruit growth.
	Fertigation	8-10L	1:100	
BROADACRE Cereals Canola Legumes	Foliar	2-3L	1:20	Apply from early emergence up to tillering. Apply again if iron deficiency exists.
	Foliar	2-3L	1:20	
	Foliar	2-3L	1:20	
CAPSICUM	Foliar	2L	1:100	Apply when sufficient leaf area is present. Repeat at fruit set and development.
	Fertigation	5-10L	1:100	
CITRUS	Foliar	2-3L	1:100	Apply as required to maintain trace element levels.
	Fertigation	10-15L	1:100	
COTTON	Aerial	2L	Use maximum practical water	Apply as required to maintain trace element levels.
	Ground rig	3-7L	1:50	
GRAPES/ VINES	Foliar	2-3L	1:100	Maximum 3 applications may be required post flowering up to veraison.
	Fertigation	10-12L	1:100	
OLIVES	Foliar	2-3L	1:100	Apply early in the season, repeat to maintain trace elements levels. Repeat post-harvest.
	Fertigation	9-15L	1:100	

CROP		RATE/ha	Minimum dilution	Comments
PISTACHIOS	Foliar	2-3L	1:100	Apply as required to maintain crop colour and vigour. Repeat mid-late fruit growth.
	Fertigation	10-15L	1:100	
POTATOES	Foliar	2-3L	1:100	Apply as required to maintain crop colour and vigour. Repeat mid-late fruit growth.
	Fertigation	9-12L	1:100	
STONE FRUIT	Fertigation only	10-15L	1:100	Apply pre-flowering or on recently hardened flush. Repeat as necessary.
STRAWBERRIES	Foliar	2L	1:100	Apply 3-4 weeks after transplanting. Repeat as required.
	Fertigation	6-10L	1:100	
SUGARCANE	Foliar	2-3L	1:100	Apply during early growth when sufficient leaf area. Apply in conjunction with billet fungicide.
	Fertigation	4-5L	1:100	
TOMATOES	Foliar	2L	1:100	Apply 3 weeks after transplanting. Repeat at first truss and fruit development.
	Fertigation	10-15L	1:100	
TURF	Foliar	30L	1:100	As required to maintain turf quality and improve colour.

ANALYSIS	w/v%	Impurities
Iron (Fe) as EDTA	6.0	Mo < 0.001% Se < 0.001% Cd < 1mg/kg Pb < 5mg/kg Hg < 0.2mg/kg

Specific Gravity 1.28 ± 0.01

**DO NOT apply to the foliage or fruit of sensitive crops such as apricots, nectarines, plums & pome fruit as phytotoxicity may occur.**

**Water Ratio:** For foliar application use 1:100 minimum water dilution rates. For Aerial application use maximum practical water rates. The rates must be adjusted according to the tissue and/or soil analysis results. Note: Only mix diluted chelate formulations with Liquid NPK's and do not store solution.



Liquid Fertilizer: Do not swallow, may be harmful. The spray from this product may act as an irritant. Avoid Inhalation and contact with eyes and skin.

**Not classified as a hazardous substance according to the criteria of Safe Work Australia.**  
**Prevention:** Wear protective gloves/protective clothing/eye protection/face protection.  
**Storage & Disposal:** Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

**SPRAYGRO LIQUID FERTILIZERS**  
 76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494  
 PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496  
 PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au  
**PRODUCED AND MANUFACTURED IN AUSTRALIA**

NOT APPLICABLE 1000 LITRE CONTAINERS