

CAUTION

03 SR-7

DISCLAIMER/CONDITIONS OF SALE: (1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and/or user of Spraygro products ("customer") in respect to any loss or damage including consequential loss or damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. (2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon any descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. (3) The customer understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. (4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. (5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. (6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.



Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

- Use the correct label spray rates of any pesticide used with Spraygro product.
- **DO NOT** use when crop is under heavy stress from pests or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **DO NOT** spray to visible droplet coalescence. **DO NOT** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is imminent.
- Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
- Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.

DRIPRO®

SMARTTRACE®

CHELATED ELEMENT

IRON

GUARANTEED MINIMUM ANALYSIS
IRON AS LIGNOSULFONATE 5.0% w/v
AND SULFUR 2.9% w/v

20 Litre NET

200 Litre NET

1000 Litre NET

Application recommendations: Spraygro Smartrace Iron is suitable for the following crops when an iron deficiency exists. Apply with sufficient water for thorough coverage of leaves. The rates are to be adjusted according to the tissue and/or soil analysis results.

BROADACRE: Rate of application per min water/hectare: 50 litres

LUCERNE	@ 1-2 litres/hectare. Apply pre-flowering or earlier if deficiency is evident.
PASTURE	@ 2-3 litres/hectare. Apply when deficiency is evident.
CEREAL	@ 2-3 litres/hectare. Apply mid-late tillering. Repeat as necessary.
COTTON	@ 2-3 litres/hectare. Apply 4-6 weeks after planting or when deficiency exists.

HORTICULTURE: Rate of application per min water/hectare: 300 litres

CITRUS	@ 3-6 litres/hectare. Apply in springtime as a thorough spray, up until bloom and again after bloom if necessary. Best absorption occurs 1/2 - 1/3 through a flush of growth. Spray should not be applied during blossoming or harvest.
POME FRUIT	@ 0.5 litres/hectare. Apply as required at fruit development.
ALMONDS	@ 3-4 litres/hectare. Apply when deficiency is evident.
GRAPES	@ 2-4 litres/hectare. Apply when deficiency is evident.
STRAWBERRIES	@ 2 litres/hectare. Apply 3-4 weeks after transplanting, repeat at fruit development.
VEGETABLES	@ 3-5 litres/hectare. Apply when plants are young and before flowering. Repeat as necessary.

WARNING: DO NOT apply to the foliage or fruit of sensitive crops such as apricots, nectarines & plums as phytotoxicity may occur.

TURF: Rate of application:

TURF	@ 100-400mls/100m ² /water dilution 1:100. Apply to enhance turf colour year round.
ORNAMENTALS	Foliar @ 2 litres/per 1000 litres water.
SHRUBS	Soil application drench @ 8 liters per 1000 litres water.

ANALYSIS	w/v%
Iron (Fe) as lignosulfonate	5.0

Specific Gravity 1.15

Impurities

Mo < 0.001%
Se < 0.001%
Cd < 1mg/kg
Pb < 1mg/kg
Hg < 0.2mg/kg

COMPATIBILITY: Highly incompatible with Amine salts of 2,4D or MCPA (excludes LVE versions).

May block filters when combined with propiconazole. It is recommended to apply non-ionic wetter (e.g., BS1000) to improve compatibility. Add propiconazole first, then wetter and then **under agitation** add this product.

Liquid Fertilizer: Do not swallow, may be harmful. The spray from this product may act as an irritant. Avoid Inhalation and contact with eyes and skin.

Not classified as a hazardous substance according to the criteria of Safe Work Australia.

Prevention: Wear protective gloves/protective clothing/eye protection/face protection.

Storage & Disposal: Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.



NOT APPLICABLE 1000 LITRE CONTAINERS

SPRAYGRO LIQUID FERTILIZERS
76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494
PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au

PRODUCED AND MANUFACTURED IN AUSTRALIA