



DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect to any loss or damage including consequential loss or damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon any descriptions, illustrations or specifications containede Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representations of the type set out above is on the basis that the customer independently verify the accuracy thereof. 4) the customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or disclaimer concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.

Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

- **DO NOT** apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is imminent.
- **DO NOT** spray to run off. **DO NOT** spray to visible droplet coalescence. **DO NOT** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** use when crop is under heavy stress from pests or inadequate moisture levels.
- Use the correct label spray rates of any pesticide used with Spraygro product.
- Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
- If precipitation occurs, do not proceed further.
- Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture.
- The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.

CAUTION

12 BS-4



BLUESTRIPE

CAUTION
 KEEP OUT OF REACH OF CHILDREN
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING

COPPER SULFATE

To be used where a Copper deficiency occurs
 GUARANTEED MINIMUM ANALYSIS

COPPER.....6.7% w/v. SULFUR.....3.4% w/v.

20 Litre NET 200 Litre NET 1000 Litre NET

BLUESTRIPE COPPER SULFATE is suitable for the following crops where a Copper deficiency occurs. Apply with sufficient water for thorough coverage of leaves. The rates must be adjusted according to tissue and or soil analysis results.

CROPS	RATE PER HECTARE	CRITICAL COMMENTS	Impurities
Cereals	1.5-3 litres/ha	Apply early-mid tillering	Mo < 0.001%
Canola	1.5-3 litres/ha	Apply pre-bolting where a deficiency occurs.	Se < 0.001%
Legumes	1.5-3 litres/ha	Apply pre-flowering or earlier if a deficiency is evident.	Cd < 1mg/kg
Pasture	1.5-3 litres/ha	Apply when a deficiency is known to exist.	Pb < 6mg/kg
Lucerne	1.5-3 litres/ha	Apply when deficiency is evident pre-flowering.	Hg < 0.2mg/kg

COMPATIBILITY: Highly incompatible with Amine salts such as 2,4D and MCPA (excludes LVE) and with propiconazole.
SPRAYING RATE: For foliar spraying through conventional spraying equipment, use a minimum of 150 litres water per hectare. When using low volume equipment, 75 litres per hectare is usually sufficient.
CROP SAFETY: Avoid spraying sulfate trace elements when conditions cause excessive evaporation and leaf burn. Be aware the addition of nitrogen fertilizer to this product may increase the risk of phytotoxicity. **DO NOT** apply to the foliage or fruit of sensitive crops such as apricots, nectarines, plums & pome fruit as phytotoxicity may occur.

WARNING
Hazards: Harmful if swallowed, Harmful in contact with skin. Causes skin and serious eye irritation. Toxic to aquatic life and harmful with long-lasting effects.
Prevention: Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Avoid release to the environment. Wear protective gloves/protective clothing/eye protection/face protection.
Response: Call a POISON CENTER or doctor if you feel unwell. Rinse mouth. Take off contaminated clothing. Wash contaminated clothing before use. IF SWALLOWED do NOT induce vomiting and Immediately call a POISON CENTER or doctor. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do – continue rinsing. If eye or skin irritation persists get medical advice/attention. For advice, contact a Poisons Information Centre (e.g. phone Australia 13 11 26; New Zealand 0800 764 766) or a doctor (at once).
Storage & Disposal: Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.



NOT APPLICABLE 1000 LITRE CONTAINERS

SPRAYGRO LIQUID FERTILIZERS
 76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494
 PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
 PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au

PRODUCED AND MANUFACTURED IN AUSTRALIA