

## CAUTION

The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
- Use the correct label spray rates of any pesticide used with Spraygro product.
- **DO NOT** use when crop is under heavy stress from pests or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **DO NOT** spray to visible droplet coalescence. **DO NOT** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is imminent.

Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

**DISCLAIMER/CONDITIONS OF SALE:** (1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect to any loss or damage including consequential loss or damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. (2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon any descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. (3) The customer understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. (4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. (5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. (6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.

21 BS-5

**Application recommendations:** Boron-15 suitable for the following crops when boron deficiency exists. Apply with sufficient water for thorough coverage of leaves. The rates are to be adjusted according to the tissue and/or soil analysis results.

**BROADACRE:** Barley, Canola, Cotton, Legumes, Maize, Rice, Wheat, Cane, Sunflower, Lucerne & Soya beans.

Foliar Rate/Ha	Fertigation/Ha	Comments
1-2L	n/a	Foliar apply in a minimum of 50 - 100L final spray volume for ground rigs or 10-20L in a minimum of 25-50L final spray volume for Aerial rigs. Apply as required prior to flowering.

**NUT TREES:** Almond, Pistachio & Walnut.

1-2L	n/a	Foliar apply in a minimum of 200 - 400L final spray volume. 3 Applications: 1st at early spur burst, 2nd at complete petal fall 3rd post harvest at 3L/ha.
------	-----	--

**FRUIT TREES:** Apple, Cherry, Nectarine, Peach, Pear & Pomegranate.

1-2L	n/a	Foliar apply in a minimum of 200 - 400L final spray volume. 3 Applications: 1st at early spur burst, 2nd at complete petal fall 3rd post harvest at 3L/ha. Note Do NOT apply as foliar to stone fruit during leaf growth, can be applied post harvest but before leaf drop.
------	-----	---

**EVERGREEN TREE CROPS:** Avocado, Citrus, Macadamia & Lychee.

1-2L	2-3 L	Foliar apply in a minimum of 200 - 400L final spray volume. Apply prior to fruit bud development, repeat at spring flush. A post harvest application may also be required.
------	-------	--

**FRUITING VEGETABLES:** Capsicum, Cucurbits, Eggplant, Tomatoes (field), Watermelon & Pumpkins.

1-2L	2-3L	Foliar apply in a minimum of 200 -400L final spray volume. Apply at 4-6 true leaf stage or as required. When practical use higher (more dilute) water rates.
------	------	--

**LEAFY VEGETABLES:** Endive, Fennel Lettuce, Broccoli, Cabbage, Cauliflower, Kale & Herbs.

Foliar Rate/Ha	Fertigation/Ha	Comments
1-2L	2-3L	Foliar apply in a minimum of 200 - 400L final spray volume. Apply as required. DO NOT APPLY IN HEAT OF DAY.

**ROOT VEGETABLES:** Beetroot, Carrot, Leek, Onion, Potato, Radish & Sweet Potato.

1-2L	2-3L	Foliar apply in a minimum of 200 - 400L final spray volume. Apply as required. DO NOT APPLY IN HEAT OF DAY.
------	------	---

**VINE and BERRY CROPS:** Blueberry, Strawberry, Raspberry, Wine & Table grapes.

1-2L	2-3L	Foliar apply in a minimum of 200 - 400L final spray volume. In season treatments 3 each at: 1st cluster visible, 2nd at flower buds separated, 3rd at fruit set. Do not exceed 1 x concentration or Ha rate. Post harvest treatment. Select either a foliar or fertigation treatment, but not both.
------	------	---

**POPPIES**

2-3L	n/a	Foliar apply in a minimum of 50 - 100L final spray volume for ground rigs or 10-20L in a minimum of 25-50L final spray volume for Aerial rigs. Apply as required prior to flowering. Apply from row over til capsule development as needed.
------	-----	---

ANALYSIS	w/v%	w/w%
Nitrogen (N)	7.5	5.5
Boron (B)	15	10.9

Specific Gravity 1.375

**NOTE:** Do not foliar apply more than 3 litres of Boron-15 per hectare in any one spray application. If crop requires more than this do it in split applications allowing 14 days between sprays. **SOIL APPLICATIONS:** Boron-15 may be applied uniformly across the field. DO NOT apply combination of Boron-15 with (mancozeb + oil) or (Dithane OC).

Liquid Fertilizer: Do not swallow, may be harmful. The spray from this product may act as an irritant. Avoid Inhalation and contact with eyes and skin.

**Not classified as a hazardous substance according to the criteria of Safe Work Australia.**

**Prevention:** Wear protective gloves/protective clothing/eye protection/face protection.

**Storage & Disposal:** Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

**SPRAYGRO LIQUID FERTILIZERS**

76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494  
PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496  
PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au

**PRODUCED AND MANUFACTURED IN AUSTRALIA**

**Impurities**

Mo < 0.001%

Se < 0.001%

Cd < 1mg/kg

Pb < 5mg/kg

Hg < 0.2mg/kg



NOT APPLICABLE 1000 LITRE CONTAINERS

20 Litre NET

200 Litre NET

1000 Litre NET

Batch No. 9

