



**HORTICULTURE: Foliar applications per min water/hectare generally required: 150-300L. Tree crops may require higher rates (500-1000L)**

CROP	FOLIAR RATES L/Ha	FERTIGATION RATES L/Ha	COMMENTS
ALMONDS	5-7L	15-30L	Apply 2 weeks post-bloom. Repeat 2-3 times every 4 weeks.
APPLES	5-7L	15-25L	Apply at petal fall. Continue at 7-14 day intervals until harvest.
ASPARAGUS	4-6L	8-12L	Apply 7 day from 1st fern. Repeat at spear emergence.
AVOCADOS	3-5L	10-15L	Apply 3 weeks prior to flowering. Repeat every 3 weeks from fruit set to harvest.
CARROTS & ONIONS	3-4L	10-12L	Apply weekly from root differentiation until 1 week before harvesting carrots. Apply from bulb development at regular fortnightly intervals in onions.
BROCCOLI	2-3L	7-15L	Apply every 2 weeks until head formation.
CAPSICUM	2-3L	9-15L	Apply every 2 weeks until flowering. Repeat at regular intervals during fruiting.
CELERY	2-3L	7-12L	Apply every 2 weeks. Stop application 1 week before harvest.
CITRUS	5-7L	10-25L	Apply prior to bloom, at petal fall and later as required.
COTTON	5-6L	10-20L	Apply prior to 1st flowering. Repeat at boll formation.
CUCUMBERS & CUCURBITS	2-3L	8-12L	Apply prior to flowering with further applications at regular intervals. Do not apply a week before harvest.
GRAPE VINES	2-3L	10-25L	Apply prior to flowering with regular applications until 2-3 weeks before harvesting.

**DO NOT apply K Blast 30 to foliage of crops sensitive (foliar burn) to sulfur.** Do not apply K Blast 30 foliar with crop oil sprays. Allow at least 14 days before or after an application of crop oil when applying K Blast 30 as a foliar. When fertigating allow greater than 1 hour injection period with other liquid fertilizers as well as apply ample irrigation water immediately before and after to ensure even distribution of K Blast 30 to the crop and prevent crop injury.

**WARNING:** Highly incompatible with low pH products (acids) and chemical reaction may occur resulting in toxic sulfur gases. Product pH is approximately 7.5. Blends of product must not be acidified below a pH of 6.0.  
Liquid Fertilizer: Do not swallow, may be harmful. The spray from this product may act as an irritant. Avoid Inhalation and contact with eyes and skin.

**Not classified as a hazardous substance according to the criteria of Safe Work Australia.**  
**Prevention:** Wear protective gloves/protective clothing/eye protection/face protection.  
**Storage & Disposal:** Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.



NOT APPLICABLE 1000 LITRE CONTAINERS

**SPRAYGRO LIQUID FERTILIZERS**  
76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494  
PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496  
PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au

**PRODUCED AND MANUFACTURED IN AUSTRALIA**

**Impurities**  
Mo < 0.001%  
Se < 0.001%  
Cd < 1mg/kg  
Pb < 1mg/kg  
Hg < 0.2mg/kg

Specific Gravity 1.46

26 NH-4

**CAUTION**

The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
- Use the correct label spray rates of any pesticide used with Spraygro product.
- **DO NOT** use when crop is under heavy stress from pests or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **DO NOT** spray to visible droplet coalescence. **DO NOT** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is imminent.

Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

**DISCLAIMER/CONDITIONS OF SALE:** 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and/or user of Spraygro products ("customer"), in respect to any loss or damage including consequential loss or damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon any descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. 4) the customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.

