41 BS-1

CAUTION

The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
- Use the correct label spray rates of any pesticide used with Spraygro product.
- **DO NOT** use when crop is under heavy stress from pests or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **DO NOT** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day when temperature exceeds 28^{o}C, high humidity situations or when rain or frost is imminent.

Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

Batch:

DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect to any loss or damage including consequential loss or damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon any descriptions, illustrations or specifications contained Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. 4) the customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred of suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.

9 333056 008999

MolyIT application recommendations: Apply to the following crops where a deficiency exists. MINIMUM DILUTION: 1:100

CROP	FOLIAR RATES mL/Ha	Minimum Dilution	COMMENTS
APPLES	300-375	1:1000	For best results apply onto new foliage.
ASPARAGUS	150-325	1:700	Apply when required.
AVOCADOS	325-625	1:1000	Apply when required or apply at 4-5 leaf stage.
BRASSICAS	325-625	1:400	Apply at 4-6 leaf stage. Follow up applications at 7-10 day intervals as required.
CAPSICUM	150-325	1:800	Apply when required or apply at 4-5 leaf stage.
CELERY	100-250	1:1000	Apply when required or apply at 4-5 leaf stage.
CITRUS	200-325	1:2000	For best results apply onto new foliage when required.
COTTON/ CEREALS	250-375	1:500	For best results apply onto new foliage when required.
CUCURBITS	325-625	1:400	Apply when required or apply at 4-5 leaf stage.
GRAPE VINES	250-450	1:1500	For best results apply onto new foliage when symptoms are exposed.
LEGUMES	325-625	1:400	Apply when early growth is approx. 150mm tall to ensure effective movement into nitrogen fixing nodules.
LETTUCE	325-625	1:400	Apply at 2-4 leaf stage. Follow up applications at 7-10 day intervals as required.
NUT CROPS	125-325	1:2000	For best results apply onto new foliage when symptoms are exposed.

CROP	FOLIAR RATES mL/Ha	Minimum Dilution	COMMENTS
OLIVES	100-325	1:2000	For best results apply onto new foliage when required.
LUCERNE*	200-325	1:800	Apply when early growth is approx. 150mm tall to ensure effective movement into nitrogen fixing nodules. Apply at least 2 weeks before cutting every third cut.
PASTURE*	65-100	1:800	Apply when early growth is approx. 150mm tall to ensure effective movement into nitrogen fixing nodules. One application per year.
POTATOES	125-325	1:1000	Apply as symptoms are exposed or apply at 4-5 leaf stage.
STRAWBERRIES	150-325	1:400	Apply when required or apply at 4-5 leaf stage.
TOMATOES	250	1:250	Apply when required or apply at 4-5 leaf stage.
TREE CROPS	325-625	1:1000	For best results apply onto new foliage when required.
VEGETABLES	150-325	1:400	Apply when required or apply at 4-5 leaf stage.
ZUCCHINI	125-325	1:400	Apply when required or apply at 4-5 leaf stage.

*__WARNING__: Excessive use of molybdenum (Mo) can be harmful to stock. Plant levels of Mo can be high for 4 weeks after application. It is advisable to keep stock off treated areas for this period. Molybdenum application may induce copper deficiency in grazing animals. This is most likely to occur on sandy soils low in copper. Check rate and frequency of molybdenum use with appropriate authorities.

The rates must be adjusted according to the tissue and/or soil analysis results.

ANALYSIS	w/v%
Molybdenum (Mo) as organic acid molybdate complex	20.0
Phosphorus (P)	0.00

Impurities
Se < 0.001%
Cd < 50mg/kg
Pb < 2mg/kg
Hg < 0.2mg/kg

This product contains heavy metal impurities. Its use may result in accumulation of cadmium in the soil. Depending on soil characteristics, irrigation water quality, plant species and variety, crop uptake of cadmium may lead to residue levels in plant and animal products in excess of the maximum level specified by the Australia New Zealand Food Standards Code.

Compatibility: MolyIT is designed to be mixed with most fertilizers even including calcium fertilizers.

Liquid Fertilizer: Do not swallow, may be harmful. The spray from this product may act as an irritant. Avoid Inhalation and contact with eyes and skin.

Not classified as a hazardous substance according to the criteria of Safe Work Australia.
Prevention: Wear protective gloves/protective clothing/eye protection/face protection.
Storage & Disposal: Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

Specific Gravity 1.50 ± 0.01
pH 7.0

drumMUSTER eligible container

NOT APPLICABLE 1000 LITRE

SPRAYGRO LIQUID FERTILIZERS
76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494
PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au

PRODUCED AND MANUFACTURED IN AUSTRALIA