

CAUTION

The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
- Use the correct label spray rates of any pesticide used with Spraygro product.
- **DO NOT** use when crop is under heavy stress from pests or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **DO NOT** spray to visible droplet coalescence. **DO NOT** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is imminent.

Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and/or user of Spraygro products ("customer") in respect to any loss or damage including consequential loss or damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon any descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. 4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.

Batch No. 9 333056 001556



LIQUID FERTILIZER

BASE™

N 14 : P 16 : K 18

+ Chelated Trace Elements + 1% potassium humate/fulvate

20 Litre NET 200 Litre NET 1000 Litre NET

43 HN-2

Base™ 14-16-18 + trace elements is designed primarily as a fertilizer to be used as a true 'base' fertilizer starter in horticulture. Apply to soil via fertigation or bandspray only to the following crops.

Apples & pears	@ 20 litres/hectare	Apply at silver tip and 2 weeks before harvest.
Asparagus	@ 10 litres/hectare	Apply at mid-fern.
Avocado	@ 15 litres/hectare	Apply at spring bud development and again after harvest in autumn.
Brassicas & Carrots	@ 15-30 litres/hectare	Apply pre-plant.
Cereal crops	@ 20-40 litres/hectare	Apply pre-plant.
Citrus	@ 10-20 litres/hectare	Apply during winter pre-blossom, full bloom and again at autumn flush.
Coffee	@ 30-40 litres/hectare	Apply soil application and again at mid-season.
Cucurbits	@ 30-50 litres/hectare	Apply pre-plant and 10-20 litres/hectare at 4-5 leaves.
Leafy vegetables	@ 30-50 litres/hectare	Apply pre-plant.
Macadamia nuts	@ 10-15 litres/hectare	Apply at spring flush.
Mango	@ 15-20 litres/hectare	Apply at dormancy and 10-15 litres/hectare during fruit swelling.
Nut trees	@ 15-20 litres/hectare	Apply at spring bud burst.
Pineapples	@ 4-6 litres/hectare	Apply at start of leaf growth early spring x 3-4 applications 4 weeks apart.
Potatoes	@ 15-20 litres/hectare	Apply at emergence/early growth.
Stone fruit	@ 15-20 litres/hectare	Apply at winter bud.
Strawberries	@ 15-20 litres/hectare	Apply 1-2 weeks after transplanting and at 10-15 litres/hectare at flowering.
Sugar cane	@ 10-15 litres/hectare	Apply at 2-3ft high and again at 4ft onwards.
Root vegetables	@ 30-50 litres/hectare	Apply pre-plant.
Table grapes - red	@ 10-15 litres/hectare	Apply at bud burst, berries forming 3mm and again at post harvest.
Tea	@ 15-20 litres/hectare	Apply late season.
Wine grapes - red	@ 10-15 litres/hectare	Apply at bud burst.

ANALYSIS	w/v%
Nitrogen (N) as urea	6.5
Nitrogen (N) as ammonium	7.2
Nitrogen (N) as amine	0.3
Nitrogen (N) as total	14.0
Phosphorus (P) and ammonium polyphosphate	8.0
Phosphorus (P) as potassium phosphate	8.0
Phosphorus (P) as total	16.0
Potassium (K) as potassium phosphate	18.0
Zinc (Zn) as EDTA	0.1
Manganese (Mn) as EDTA	0.08
Copper (Cu) as EDTA	0.04
Iron (Fe) as EDTA	0.1
Boron (B) as borate	0.03
Molybdenum (Mo) as molybdate	0.02
Potassium humate/fulvate	1.0

Specific Gravity 1.46 ± 0.01 pH 6-8

Liquid Fertilizer: Do not swallow, may be harmful. The spray from this product may act as an irritant. Avoid Inhalation and contact with eyes and skin.

Not classified as a hazardous substance according to the criteria of Safe Work Australia.

Prevention: Wear protective gloves/protective clothing/eye protection/face protection.

Storage & Disposal: Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

SPRAYGRO LIQUID FERTILIZERS

76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494
PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au

PRODUCED AND MANUFACTURED IN AUSTRALIA

Impurities

Se < 0.001%
Cd < 1mg/kg
Pb < 1mg/kg
Hg < 0.2mg/kg
F < 40g/kg P
Biuret < 9g/kg N



NOT APPLICABLE 1000 LITRE CONTAINERS