

concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor. directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction 9 warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with incurred of suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro customer shall at all times indemnify, and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. 4) the any descriptions, illustrations or specifications contained Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro be under no liability to the purchaser and or user of Spraygro products ("customer") in respect to any loss or DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall

test area should be sprayed and observed prior to the total crop spray. Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small

imminent. e Do Not apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is

fruit or leaves. • DO NOT spray to run off. DO NOT spray to visible droplet coalescence. DO NOT allow concentrated spray mist to run on

• DO NOT apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition. • DO NOT use when crop is under heavy stress from pests or inadequate moisture levels.

Use the correct label spray rates of any pesticide used with Spraygro product.

Spraygro product.

• Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding (fall out) in mixture. If precipitation occurs, do not proceed further.

parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter • Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and may result in damage to the plant.

The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions

**NOITUAD** 46 SRN-3

## Contains W Smart Rele ase Nitrogen<sup>™</sup> from Methylene Controlled NOT APPLICABLE TO 1000 LITRE CONTAINERS 1000 Litre



**Application rates for Balance-15** 

HORTICULTURE: Foliar applications per min water/hectare generally required: 150-300L

CROP	FOLIAR RATES L/Ha	FERTIGATION RATES L/Ha	COMMENTS  Apply at 7-14 day intervals from fruit set to harvest.		
AVOCADOES	4L	15L			
BERRIES	7-10L	20-30L	Apply early bloom/fruit set. Repeat 14-21 days.		
BRASSICA'S	4L	10-15L	Apply every 14 days from early head development until harvest.		
CAPSICUMS	4-6L	20-30L	Apply at 7-14 days from fruit set through to harvest.		
CUCURBITS	2-3L	5-10L	Apply every 7-14 days from fruit set to harvest.		
FRUIT CROPS	5-10L	20-30L	Apply from early bloom through to fruit set. Repeat at 30 days. Post harvest apply 20L/ha foliar.		
GARLIC	3-4L	12-15L	Apply from when sufficient leaf area exists to intercep spray and from bulb development to harvest.		
GRAPES	3-5L	10-15L	Apply before flowering then at regular intervals.		
LEAFY GREENS	5L	8-12L	Apply every 7-14 days from mid-crop to harvest.		
OLIVE TREES	10L	20-30L	Apply every 30 days from early to late fruiting.		
ONIONS	3-4L	12-15L	Apply from when sufficient leaf area exists to intercept spray and from bulb development to harvest.		
PEANUTS	N/A	10-20L	Apply at planting.		
POTATOES	4-6L	5-10L	Apply every 14-30 days from tuber initiation to harvest.		
STONE FRUIT	2-4L	5-10L	Apply at 4-6 weeks prior to harvest for improved fruit sizing, firming. Post harvest apply 20L/ha.		
TROPICAL FRUIT	4L	15L	Apply every 7-14 days from fruit set to harvest.		
VEGETABLES	4-6L	12-15L	Apply every 14-21 days from mid-crop to harvest		

BROADACRE: Foliar rate of application per min water/hectare: 80 litres.

CROP	COMMENTS
COTTON	Apply 6-8L/ha at flowering an boll set. Repeat as necessary through to maturation.
LEGUME GRAIN	Apply 4L/ha during pod development.

**TURF:** Can be applied with low water rates as a foliar or use higher water rates to drench product in. Smart Release Nitrogen can release over 20-30 days on leaf surface or up to 8-10 weeks in the soil. Smart Release Nitrogen enhances the uptake of other nutrients

Apply the following rates at 4-6 week intervals for optimum turf growth.

CROP	COMMENTS				
TURF	Low feed: Medium feed: High feed:	Apply 80L/ha apply min 1:25 water dilution. Apply 120L/ha apply min 1:25 water dilution. Apply 150L/ha apply min 1:25 water dilution.			

Should you require a higher Nitrogen amount, tank mix with Smart Release Nitrogen™ Nitro-35.

ANALYSIS	w/v%	ANALYSIS	w/v%	Impurities
Nitrogen (N) as polymethylene urea	5.3	Manganese (Mn) as EDTA	0.03	Mo < 0.001%
Nitrogen (N) as total	15	Copper (Cu) as EDTA	0.02	Se < 0.001%
Phosphorus (P) as phosphate	4	Iron (Fe) as EDTA	$\begin{array}{c c} \hline 0.046 & Cd < 1mg/kg \\ Pb < 1mg/kg \end{array}$	Pb < 1mg/kg
Potassium (K)	8	Boron (B) as borate	0.04	Hg < 0.2mg/kg biruet < 9g/Kg N
Magnesium (Mg) as EDTA	0.02	Molybdenum (Mo) as molybdate	0.0016	
Zinc (Zn) as EDTA	0.03	Cobalt (Co) as EDTA	0.0014	

Liquid Fertilizer: Do not swallow, may be harmful. The spray from this product may act as an irritant. Avoid Inhalation and contact with eyes and skin.

Specific Gravity 1.25 ± 0.01 pH 10-11

**SPRAYGRO LIQUID FERTILIZERS** 

76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494 PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496 PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au

Australia. Storage & Disposal: Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.

This product is not classified as Hazardous according to the criteria of Safe Work

PRODUCED AND MANUFACTURED IN AUSTRALIA