



BULLKELP PLUS ++ is suitable for application by Boom Spray or Fertigation. Recommended minimum water dilution of 1:300.

BROADACRE: Barley, Canola, Cotton, Legumes, Maize, Oats, Rice, Sorghum, Triticale, Wheat, Sugar Cane & Pasture Crops.

| Foliar Rate/Ha | Comments |
|----------------|--|
| 5-7L | Foliar: Apply in a minimum of 600-900L final spray volume with a minimum 1:300L water dilution. Apply every 14-21 days from 5th visible leaf onwards. For best results apply first sprays before leaf hardening of crop. Apply to sugarcane at planting and during the lead up to hot, dry conditions. |

NUT TREES: Almond, Pistachio & Walnut.

| | |
|-------|--|
| 5-10L | Foliar apply in a maximum of 1000L final spray volume. Dip Rate: New Planting: 1:100 dilution. Dip bare roots for 5 minutes before transplant, or Soak seedling bag before transplant, or Soak soil around trees after plant-out. Spray at 30% bloom and repeat twice at 10 – 12 day intervals. Spray at fruit set (10 mm fruit diameter). Repeat twice at 14 day intervals. |
|-------|--|

FRUIT TREES: Apple, Cherry, Nectarine, Peach, Pear & Pomegranate.

| | |
|-------|---|
| 5-10L | Foliar apply in a maximum of 1000L final spray volume. Dip Rate: New Planting: 1:100 dilution. Dip bare roots for 5 minutes before transplant, or Soak seedling bag before transplant, or Soak soil around trees after plant-out. Spray at 30% bloom and repeat twice at 10 – 12 day intervals. Spray at fruit set (10 mm fruit diameter). Repeat twice at 14 day intervals |
|-------|---|

EVERGREEN TREE CROPS: Avocado, Citrus, Macadamia & Lychee.

| | |
|-------|--|
| 5-10L | Foliar apply in a maximum of 1000L final spray volume. Dip Rate: New Planting: 1:200-300 dilution. Spray avocado trees at the beginning of spring growth and repeat at 50% flowering. Spray macadamia trees at start of bloom and repeat 4 times at monthly intervals. Spray at fruit set (10 mm fruit diameter). Repeat twice at 14 day intervals. If water volume exceeds 1000 L/ha, increase Kelpit rate to retain 1:300 dilution. Dip bare roots for 5 minutes before transplant, or Soak seedling bag before transplant, or Soak soil around trees after plant-out. |
|-------|--|

FRUITING VEGETABLES: Capsicum, Cucurbits, Eggplant, Tomatoes (field), Watermelon & Pumpkins.

| Foliar Rate/Ha | Comments |
|----------------|--|
| 5-10L | Foliar apply in a maximum of 1000L final spray volume. Dip Rate: 1:100 dilution. Dip tray with seedlings in solution, or Wet seedling tray with watering can before transplant. Spray 14 days after transplant. Repeat once or twice at 14 day intervals. Direct seeded plants receive first foliar application at the 4-leaf stage. |

LEAFY VEGETABLES: Endive, Fennel Lettuce, Broccoli, Cabbage, Cauliflower, Kale & Herbs.

| | |
|-------|---|
| 5-10L | Foliar apply in a maximum of 800L final spray volume. Dip Rate: 1:100 dilution. Dip tray with seedlings in solution, or Wet seedling tray with watering can before transplant, or for direct seeding, Spray at 3 to 5-leaf stage. Repeat once or twice at 14 to 21 day intervals. |
|-------|---|

ROOT VEGETABLES: Beetroot, Carrot, Leek, Onion, Potato, Radish & Sweet Potato.

| | |
|-------|--|
| 5-10L | Foliar apply in a maximum of 800L final spray volume. Dip Rate: 1:300 dilution. Dip seed potatoes before planting for approximately 5 minutes, or Spray in furrow before covering, then spray after emergence at 15 cm crown diameter (14 to 28 days DAE). For other crops: Spray 14 to 21 days after transplant, or spray at 3 to 4-leaf stage for direct seeded plants. Repeat foliar application twice at 14 to 21 day intervals. Do not apply to onions or garlic after bulb swelling. |
|-------|--|

VINE and BERRY CROPS: Blueberry, Strawberry, Raspberry, Wine & Table grapes.

| | |
|-------|---|
| 5-10L | Foliar apply in a maximum of 1000L final spray volume. Dip Rate: 1:100 dilution. Spray after full bloom. Repeat 2 to 3 times at 10 to 12 day intervals (increase Kelpit rate to maintain 1:300 dilution when high water volumes are used). Apply 2 to 3 times as bunch Dips, or directed bunch Sprays at 4 to 14 mm berry size. |
|-------|---|

| ANALYSIS | w/v% | Impurities |
|------------------------------------|------|------------------|
| Nitrogen (N) as polymethylene urea | 1.2 | Mo < 0.001% |
| Total Nitrogen (N) | 5.0 | Se < 0.001% |
| Potassium (K) | 1.5 | Cd < 1mg/kg |
| | | Pb < 10mg/kg |
| | | Hg < 0.2mg/kg |
| | | biuret <15g/Kg N |

Extracts of Bull Kelp contains Auxins, Cytokinins, Alginates, Ascorbic acid, Vitamins B1, B2, C, E, Niacin, Choline, Iodine and trace elements. Specific Gravity 1.07 ± 0.01



Liquid Fertilizer: Do not swallow, may be harmful. The spray from this product may act as an irritant. Avoid Inhalation and contact with eyes and skin.

Not classified as a hazardous substance according to the criteria of Safe Work Australia.
Prevention: Wear protective gloves/protective clothing/eye protection/face protection.
Storage & Disposal: Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.



SPRAYGRO LIQUID FERTILIZERS
 76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494
 PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
 PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au
PRODUCED AND MANUFACTURED IN AUSTRALIA



CAUTION: The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.
 • Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
 • Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
 • Use the correct label spray rates of any pesticide used with Spraygro product.
 • **DO NOT** use when crop is under heavy stress from pests or inadequate moisture levels.
 • **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
 • **DO NOT** spray to run off. **DO NOT** spray to visible droplet coalescence. **DO NOT** allow concentrated spray mist to run on fruit or leaves.
 • **DO NOT** apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is imminent.
 Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

DISCLAIMER/CONDITIONS OF SALE: (1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect to any loss or damage including consequential loss or damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. (2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon any descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. (3) The customer understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. (4) The customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. (5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. (6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.

47 GP-5