

81 GR-5

CAUTION

The following conditions must be observed to apply Spraygro product blends successfully. Failure to follow these instructions may result in damage to the plant.

- Use Spraygro product under conditions of optimum plant growth including moderate humidity, moderate temperatures and adequate soil moisture. Early morning or late afternoon spraying is most effective as evaporation can occur during hotter parts of the day. When mixing with other chemicals, always test a small quantity (e.g., bucket test) and check for precipitate (fall out) in mixture. If precipitation occurs, do not proceed further.
- Add wettable or soluble powders, emulsifiers or flowables to water in the mix and wet, dissolve or disperse before adding Spraygro product.
- Use the correct label spray rates of any pesticide used with Spraygro product.
- **DO NOT** use when crop is under heavy stress from pests or inadequate moisture levels.
- **DO NOT** apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.
- **DO NOT** spray to run off. **DO NOT** spray to visible droplet coalescence. **DO NOT** allow concentrated spray mist to run on fruit or leaves.
- **DO NOT** apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is imminent.

Note: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray.

DISCLAIMER/CONDITIONS OF SALE: (1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall be under no liability to the purchaser and or user of Spraygro products ("customer") in respect to any loss or damage including consequential loss or damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro Product Profile or not. (2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon any descriptions, illustrations or specifications contained in Spraygro's Product Profile or any other document produced by Spraygro. (3) The customer understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. (4) the customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability incurred or suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. (5) Spraygro warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or direction concerning this product. (6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.

Batch No. 333056 001952





Gripper™
 edta chelate
6% COPPER w/v
 20 Litre NET 200 Litre NET 1000 Litre NET

Application recommendations: Gripper Copper is suitable for the following crops where Copper deficiency exists. Apply with sufficient water for thorough leaf coverage.

CROP	RATE/ha	Minimum dilution	Comments
ALMONDS NUT TREES	Foliar Fertigation 1-2L 4-6L	1:100 1:100	Apply as required to maintain crop colour and vigour. Repeat early fruit growth.
BROADACRE CROPS Cereals Canola	Foliar Foliar 0.5-2L	1:40	Apply to actively growing plants when sufficient leaf area exists to spray, repeat as necessary. Apply between mid-late tillering to flag leaf emergence (fle) stage. After fle apply maximum of 1.0L/ha with minimum of 75L water dilution. After head emergence, apply maximum of 0.5L/ha with minimum 75L water.
Legumes	Foliar		Apply at 4-8 leaf stage prior to pod set.
VEGETABLES	Foliar Fertigation 1-2L 2-4L	1:100 1:100	Apply when sufficient leaf area is present before flowering.
CITRUS	Foliar Fertigation 0.5-1L 2-4L	1:100 1:100	Apply when deficient. Use lower rates on trees up to 6 year stage.
COTTON	Aerial Ground Water 0.83-2L	Use maximum practical water 1:50	Apply prior squaring.
GRAPES/ VINES	Foliar Fertigation 0.5-1L 2-3L	1:100 1:100	Apply when deficiency exists.

CROP	RATE/ha	Minimum dilution	Comments
ONIONS	Foliar Fertigation 1L 2L	1:100 1:100	Apply at 2-3 leaf stage Apply at bulb formation.
POME FRUIT	Foliar Fertigation 0.5-1.0L 4-6L	1:300 1:300	Apply in Spring up until bloom. Do not apply during blossom or harvest.
POTATOES	Foliar Fertigation 1-2L 2-4L	1:100 1:100	Apply to vines when sufficient leaf area. Repeat at tuber development.
STONE FRUIT	Fertigation only 3-5L	1:100	Apply pre-flowering or on recently hardened flush. Repeat as necessary.
STRAWBERRIES	Foliar Fertigation 0.5L-1L 1-2L	1:100 1:100	Apply 3-4 weeks after transplanting. Repeat at fruit development.
TOMATOES	Foliar Fertigation 2-3L 4-5L	1:100 1:100	Apply 3 weeks after transplanting. Repeat at first truss and fruit development.
TURF	Foliar 100ml/ 100m²	1:100	As required to maintain turf quality and improves colour.

ANALYSIS	w/v%
Copper (Cu) as EDTA	6.0

Specific Gravity 1.26

Impurities
 Mo < 0.001%
 Se < 0.001%
 Cd < 1mg/kg
 Pb < 3mg/kg
 Hg < 0.2mg/kg

DO NOT apply to the foliage or fruit of sensitive crops such as apricots, nectarines, plums & pome fruit as phytotoxicity may occur. Be aware the addition of nitrogen fertilizer to this product may increase the risk of phytotoxicity.
Water Ratio: For foliar application use 1:100 minimum water dilution rates. For Aerial application use maximum practical water rates. The rates must be adjusted according to the tissue and/or soil analysis results. Note: Only mix diluted chelate formulations with Liquid NPK's and do not store solution.

WARNING:
Hazards: Harmful if swallowed. Harmful in contact with skin. Causes serious eye irritation.
Prevention: Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Wear protective gloves/protective clothing/eye protection/face protection.
Response: Call a POISON CENTER or doctor if you feel unwell. Rinse mouth. Wash contaminated clothing before reuse. IF SWALLOWED: Immediately call a POISON CENTER or doctor. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do – continue rinsing. If eye irritation persists get medical advice/attention.
Storage & Disposal: Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the empty container according to approved local practices, contact primary industries for disposal information.



NOT APPLICABLE 1000 LITRE CONTAINERS

SPRAYGRO LIQUID FERTILIZERS
 76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494
 PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496
 PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au
PRODUCED AND MANUFACTURED IN AUSTRALIA