

limiting trace element.

ALMONDS

PISTACHIO

VEGETABLES

BROADACRE

Cereals, Canola,

Legumes, Cotton

LEGUMES, PEAS,

store solution.

PINEAPPLES

CITRUS, GRAPES, | Foliar

CROPS

BEANS

OLIVES

CROP

Foliar

Foliar

Foliar

Fertigation

Fertigation

Fertigation

Fertigation

Fertigation

Fertigation

Foliar

Foliar

PORT ADELAIDE, SOUTH AUSTRALIA 5015 • A.B.N. 47 007 974 496 PRODUCED AND MANUFACTURED IN AUSTRALIA

Storage & Disposal: Do not contaminate water, food or feed by storage or disposal. Triple rinse containers, empty rinsate into spray tank. Dispose of the

76-80 GRAND TRUNKWAY, GILLMAN, SOUTH AUSTRALIA 5013 • PO BOX 494 PHONE: (08) 8447 7266 • Email: customerservice@spraygro.com.au www.spraygro.com.au

SPRAYGRO LIQUID FERTILIZERS

fruit or leaves.

• DO NOI use when crop is under heavy stress from pests or inadequate moisture levels.

test area should be sprayed and observed prior to the total crop spray.

NOITUAD

Specific Gravity 1.28 ± 0.01

Impurities

Mo < 0.001%

Se < 0.001%

Cd < 1mg/kg

Pb < 5mg/kg

Hg < 0.2mg/kg

pH 8.4 to 9.4

w/v%

6.8

5.0

5.0

0.23

direction concerning this product. 6) Before using please obtain a current Product Profile which is available from Spraygro or our distributor.

directions under normal conditions of use. No one other than an authorised officer of Spraygro is authorised to make any warranty, guarantee or

understands that any representation of the type set out above is on the basis that the customer independently verify the accuracy thereof. 4) the

warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with

incurred of suffered by Spraygro arising from any proceedings or threatened proceedings against Spraygro by or involving the customer. 5) Spraygro

any descriptions, illustrations or specifications contained Spraygro's Product Profile or any other document produced by Spraygro. 3) The customer

customer shall at all times indemnify and hold harmless Spraygro from and against all loss (including reasonable legal costs and expenses) or liability

Product Profile or not. 2) The customer warrants that there has been no reliance by the customer upon any representation made by Spraygro or upon

damage which may be suffered incurred or arise directly or indirectly in respect of such products, or the use thereof, in accordance with the Spraygro

be under no liability to the purchaser and or user of Spraygro products ("customer") in respect to any loss or damage including consequential loss or

DISCLAIMER/CONDITIONS OF SALE: 1) Foliar Fertilizers Pty Ltd trading as Spraygro Liquid Fertilizers and its servants and agents ("Spraygro") shall

• DO NOT apply by aircraft if surface wind is greater than 8 kph to assure adequate crop coverage and droplet deposition.

• DO NOT spray to run off. DO NOT spray to visible droplet coalescence. DO NOT allow concentrated spray mist to run on

e Do Not apply during the heat of the day when temperature exceeds 28°C, high humidity situations or when rain or frost is

Uote: We recommend that when applying to a crop or area for the first time, or in combination with other chemicals, a small .Jn9nimmi

897400 II 990EEE II 6

88 GK-2

Batch No.

must be adjusted according to the tissue and/or soil analysis results. Note: Only mix diluted chelate formulations with Liquid NPK's and do not Liquid Fertilizer: Do not swallow, may be harmful. The spray from this product may act as an irritant. Avoid Inhalation and contact with eyes and skin.

ANALYSIS

Nitrogen (N) as amine

Zinc (Zn) as EDTA

Boron (B) as Borate

Molybdenum (Mo) as molybdate

empty container according to approved local practices, contact primary industries for disposal information.

Not classified as a hazardous substance according to the criteria of Safe Work Australia.

Application recommendations: Gripper BZM 5:5 is suitable for the following crops where a multi trace element (Zn, B, Mo) deficiency exists based on the minimum amount of the most

Comments

Apply 1st early budding, 2nd nut development, 3rd

Apply to early vegetative growth to correct trace

Apply from early emergence up to tillering, stem

elongation, prior flag leaf emergence in cereals.

Apply from early emergence where there is high

Apply to early vegetative growth to correct trace

Apply early in the season, repeat to maintain trace

Water Ratio: For foliar application use 1:100 minimum water dilution rates. For Aerial application use maximum practical water rates. The rates

Prevention: Wear protective gloves/protective clothing/eye protection/face protection.

Apply as required to maintain trace element

elements levels. Repeat post-harvest.

14 days later. Water rate min 1000L/ha.

element deficiencies.

zinc requirement.

element deficiencies.

DO NOT apply to the foliage or fruit of sensitive crops such as apricots, nectarines, plums & pome fruit as phytotoxicity may occur.

levels.

Apply with sufficient water for thorough leaf coverage.

Minimum

dilution

1:100

1:100

1:100

1:100

1:100

1:100

1:100

1:100

1:100

1:100

1:100

1:100

RATE/

ha

1.5-3L

3-6L

2-4L

5-10L

2-3L

4-6L

3-5L

6-10L

1-2L

3-5L

2-3L

6-10L